



Palos Verdes Kelp Forest Restoration Project

Project Year 8: July 2020 – June 2021 (SC-3108)

Prepared for: California Department of Fish and Wildlife

Prepared by:

The Bay Foundation

Rilee Sanders, Ben Grime, Heather Burdick, Tom Ford

Vantuna Research Group

Jonathan Williams, Chelsea Williams, Dan Pondella

November 2021



The Bay Foundation
8117 W. Manchester Ave. #750
Playa Del Rey, CA 90045
(888) 301-2527
www.santamonicabay.org

3. Report of Kelp Restoration Activities Including Stated Components in Scientific Collecting Permit (SCP).

A) Kelp Restoration Goals

The *Macrocystis pyrifera* (giant kelp) canopy cover at Palos Verdes Peninsula has decreased by approximately 80% since the first large-scale survey in 1911 (Ford and Meux 2010, MBC 2019). Sedimentation, development, urban runoff and storms slowed kelp growth. At the same time, the loss of key urchin predators and competitors allowed urchins to overrun the reef and devour the remaining kelp. Subtidal observations based upon mapping efforts conducted in 2010 identified large expanses of nearshore rocky reef that were dominated by high densities of *Strongylocentrotus purpuratus* (purple sea urchins) and *Mesocentrotus franciscanus* (red sea urchins). In total, 152 acres were described to exist in an urchin barren state.

It is within this context that The Bay Foundation initiated the Palos Verdes Kelp Restoration Project through *in situ* culling of *S. purpuratus* on the Palos Verdes Peninsula. The goal is to reduce populations of *S. purpuratus* to natural densities (associated with stable giant kelp communities in southern California) to catalyze recruitment and development of giant kelp and other macroalgae. Decreased *S. purpuratus* grazing pressure allows for the enhancement of the biogenic habitat of rocky reefs that have historically supported kelp forests. Ultimately, this increases the spatial and temporal stability, as well as biomass and production associated with the kelp forests/rocky reefs on the Palos Verdes Peninsula.

B) Timeline of Restoration Goals

Restoration and monitoring activities have been conducted in kelp reference, restoration and barren sites since July 2013. The field work involved in this project is subject to sea state, oceanographic conditions, and weather. Urchin suppression efforts have expanded each year to encompass two coves (Underwater Arch and Honeymoon), and three open shore areas (Marguerite, Resort Point, and Hawthorne). These areas are located somewhat centrally on the Palos Verdes Peninsula. These sites are nearly contiguous and share similarities in ocean exposure. An additional site, Point Fermin, was started to the south and east of these other locales in the summer of 2015. Point Fermin is roughly the south-east terminus of the Palos Verdes Peninsula. White Point was established as a new site in summer 2018. Pre-restoration monitoring at this site in Year 6 described mean *S. purpuratus* densities over a 1.8-acre expanse at 67.8 urchins per m². Monitoring and additional surveys of the barren area at White Point have continued beyond the end of the Year 7 reporting deadline, resulting in expanding the total barren area to 15 acres. The unrestored portions of this site are primarily devoid of fleshy macroalgae, while the substrate is dominated by crustose coralline algae, bare rock and *S. purpuratus*. During this reporting period of July 1, 2020 through June 30, 2021 (Year 8) of the project, all restoration efforts were focused at White Point.

The progression of restoration activities is outlined in Table 1, while Table 2 provides hours of diver effort to achieve these results. Restoration efforts projected for this operational year, July 1, 2020 through June 30, 2021, are listed in Table 3.

Table 1. Restoration progress by site Years 1 through 8. Marguerite includes Marguerite North, South and Central. Specific areas restored at Underwater Arch Cove in Years 1 and 2 were re-cleared in Years 4 and 5 due to infiltration from an assumed *S. purpuratus* refuge population in a large and shallow tide pool.

| Site Name | Area Cleared (Acres) Year 1 July 2013 - June 2014 | Area Cleared (Acres) Year 2 July 2014 - June 2015 | Area Cleared (Acres) Year 3 July 2015 - June 2016 | Area Cleared (Acres) Year 4 July 2016 - June 2017 | Area Cleared (Acres) Year 5 July 2017- June 2018 | Area Cleared (Acres) Year 6 July 2018- June 2019 | Area Cleared (Acres) Year 7 July 2019- June 2020 | Area Cleared (Acres) Year 8 July 2020- June 2021 | Total Area (acres) |
|----------------------|--|--|--|--|---|---|---|---|--------------------|
| Honeymoon Cove | 4.84 | 3.56 | 0 | 0 | 0 | 0 | 0 | 0 | 8.40 |
| Underwater Arch Cove | 3.77 | 4.49 | 0 | 2.34 | 0.28 | 0 | 0 | 0 | 10.88 |
| Marguerite | 0 | 5.07 | 3.68 | 5.27 | 0 | 0 | 0 | 0 | 14.01 |
| Hawthorne | 0 | 2.72 | 1.56 | 0 | 0.89 | 0 | 0 | 0 | 5.17 |
| Point Fermin | 0 | 0 | 3.93 | 1.13 | 0.22 | 0 | 0 | 0 | 5.28 |
| Resort Point | 0 | 0 | 0 | 0 | 3.78 | 0.22 | 0 | 0 | 4.00 |
| White Point | 0 | 0 | 0 | 0 | 0 | 3.11 | 4.38 | 1.33 | 8.82 |
| Total Area | 8.61 | 15.84 | 9.16 | 8.74 | 5.17 | 3.33 | 4.38 | 1.33 | 56.55 |

Table 2. Total diving effort towards project goals July 1, 2013 through June 30, 2021.

| July 1 2013 - June 30 2021 | | |
|----------------------------------|-----------------|-------------|
| Effort (dive hours) | Monitoring | Restoration |
| The Bay Foundation | 1923.90 | 71.93 |
| Commercial Sea Urchin Harvesters | 0 | 7080.83 |
| LA Waterkeeper | 133.37 | 1030.86 |
| Subtotal | 2057.27 | 8183.62 |
| Total Dive Hours | 10240.89 | |

Table 3. Restoration areas targeted for July 1, 2020 through June 30, 2021. Restoration work will target the site listed in the table below. Periodic monitoring of all sites will continue to ensure that *S. purpuratus* densities remain at no more than two per square meter. All sites are monitored with the following methods: video transects, photo points, urchin dissections, and response monitoring. Exploration of rocky reef along Palos Verdes will continue to identify existing or potentially emergent urchin barrens in the coming year.

| Site Name | Estimated Total Barren Area (Acres) | Start Date | Total Restored Area (Acres) | Area Restored 7.1.2020-6.30.2021 (Acres) | Status | Centroid |
|-------------|-------------------------------------|---------------|-----------------------------|--|-------------|------------------|
| White Point | 15 | July 10, 2018 | 8.82 | 1.33 | In progress | 33.713, -118.315 |

Table 4. Restoration start and completion dates for all sites. Dates are based on TBF biologist post monitoring dates for each site.

| Site Name | Post Restoration Started | Restoration Completed | Notes *start/completion date based on post monitoring date |
|----------------------|--------------------------|-----------------------|--|
| Honeymoon Cove | 11/4/13 | 1/6/15 | Constant work, no inactive periods |
| Underwater Arch Cove | 7/31/13 | 1/6/15 | Main restoration accounting for 8.26 acres; Intrusion from tidepool requiring additional clearing in 2.62 acres from 4/7/17 - 7/6/17. |
| Marguerite | 10/2/15 | 6/23/17 | 6 month break from 11/24/15 - 6/27/16 on account of wasting disease |
| Hawthorne | 1/20/15 | 5/31/16 | 14 month break from 5/31/16 to 7/25/17 where 0.89 acres were restored ending work on 8/25/17 |
| Point Fermin | 7/22/15 | 7/7/17 | Initial work from 7/22/15 through 2/4/16; 7 month break until 10/7/16 where work continued til 12/14/16; then 7 month break until 0.22 acres on 7/7/17; subsequent surveys have identified large expansive barren thought to be a result from intrusion from a refuge urchin population, funding dependent still in progress |
| Resort Point | 9/20/17 | 7/3/18 | Constant work, no inactive periods |
| White Point | 7/10/18 | In Progress | Inactive periods were common during year 8 due to a combination of COVID restrictions and poor ocean conditions precluding restoration activities from occurring. The only post-restoration monitoring occurred from 9/15/20 through 11/11/20. Restoration activities have begun again during the year 9 reporting period and are ongoing. |

C) Description of Restoration, Control, and Reference Sites

All project restoration and reference sites are located on the Palos Verdes Peninsula, Los Angeles County, California. Table 5 (below) shows all potential restoration sites along with the area in hectares initially described in 2010 surveys, and representative central GPS coordinates for each.

Table 5. Area and GPS coordinates for restoration, reference and control sites.

| Restoration Site Name | Area (Hectares) | Perimeter (Meters) | Centroid (Decimal Degrees) |
|--|------------------------|---------------------------|-----------------------------------|
| Honeymoon Cove | 4.07 | 1,509 | 33.764, -118.423 |
| Christmas Tree Cove | 4.09 | 2,264 | 33.761, -118.419 |
| Marguerite | 5.19 | 2,522 | 33.757, -118.418 |
| Underwater Arch | 5.36 | 2,183 | 33.752, -118.415 |
| Hawthorne | 8.96 | 1,789 | 33.747, -118.414 |
| Portuguese Point | 1.73 | 1,604 | 33.737, -118.376 |
| Inspiration Point | 2.57 | 1,965 | 33.736, -118.368 |
| White Point | 6.07 | 2,395 | 33.713, -118.315 |
| Point Fermin | 4.37 | 3,367 | 33.704, -118.291 |
| The following sites were identified as urchin barrens in 2010 and are located within the Marine Protected Areas (MPAs) surrounding Point Vicente. Thus far these sites have only been monitored and will continue to be monitored as part of the experimental design of the overall project. Three of these sites received restoration work in the past, pre-MPA, (2005-2011) i.e., Kaplan Cove, Long Point and Old Marineland. Restoration work was conducted on a limited basis inside the MPA in the early part of 2012. Further restoration efforts within the MPAs might yield benefits to the goals of the MPAs generally and specifically to the MPA cluster on PV. | | | |
| Site Name | Area (Hectares) | Perimeter (Meters) | Centroid (Decimal Degrees) |
| Point Vicente East | 4.8 | 2,812 | 33.740, -118.406 |
| Kaplan Cove | 2.3 | 1,115 | 33.737, -118.401 |
| Long Point | 0.82 | 1,240 | 33.736, -118.398 |
| Old Marineland | 1.2 | 744 | 33.737, -118.395 |
| 120 Reef | 1.74 | 1,226 | 33.738, -118.392 |
| Abalone Cove Kelp | 9.1 | 3,397 | 33.740, -118.385 |
| Reference Site Name | Area (Hectares) | Perimeter (Meters) | Centroid (Decimal Degrees) |
| Point Vicente West | - | - | 33.740, -118.412 |
| Rocky Point North | - | - | 33.779, -118.426 |
| Ridges North | - | - | 33.787, -118.420 |
| Control Site Name | | | |
| Abalone Cove West | 9.10 | 3,397 | 33.740, -118.385 |
| Marguerite Central* | 5.19 | 2,522 | 33.757, -118.418 |
| *Marguerite Central started as a control site but switched to a restoration site in 2015. | | | |

D) Pre-Restoration Monitoring

Seven restoration sites have been established off Palos Verdes: Honeymoon Cove, Marguerite, Underwater Arch Cove, Hawthorne, Resort Point (a geographical extension of Honeymoon Cove), White Point, and Point Fermin. Pre-monitoring began at White Point (the current restoration site) in February 2018, although restoration activities did not commence until July 2018. Pre-restoration monitoring is conducted on all sites per California Department of Fish and Wildlife (CDFW) standards stipulated in the terms of the SCP. Restoration sites are divided into 30m by 30m blocks each comprised of 15 transects (2m by 30m swath) monitored by divers. Each 30m transect is divided into 10m long segments to estimate the density of *S. purpuratus*, *M. franciscanus*, *M. pyrifera* and a characterization of the substrate and relief. In certain instances, these blocks, or the individual transects comprising them, are truncated to fit the natural topography. This fine scale and spatially comprehensive methodology allows for greater resolution of inter-block variability and has been beneficial to the adaptive management of restoration teams. During the initial phase of the project (July 2013 to March 2014), all 15 transects (per block), covering 100% of the restoration block were pre-monitored. Field staff engaged in the adaptive management of the project noted the time-consuming nature of pre-monitoring transects in comparison to post monitoring. To continue to make progress in a manner consistent with contracts and the ecology of the region; program management staff at TBF, in consultation with National Oceanic and Atmospheric Administration (NOAA) biologists, conducted an applied power analysis on the pre-monitoring data set from July 2013 through February 2014. This analysis described no loss in statistical strength, and equally, no gain in accuracy in continuing to pre-monitor all transects within any given restoration block. Based on the applied power analysis, a reduction of sampling area by 66% allowed for a substantial increase in restoration efforts, while making the pre-restoration monitoring more efficient and cost-effective. TBF biologists pre-monitor five transects per restoration block.

The pre-restoration site map (Figure 1) is derived from data collected along the five 2m x 30m swaths per restoration block. The values of those data are extended and applied to the adjacent transects representing 6 x 30m swaths to estimate the total abundance of *M. franciscanus* and *S. purpuratus* pre-restoration and display the full block area on the maps. All data collected (i.e. date, area, team members, level of effort, *M. franciscanus* and *S. purpuratus* densities, *M. pyrifera* density, rugosity, and substrate) are entered, quality assured and quality controlled (QAQC), and managed utilizing a georeferenced database.

During Year 8 of the project, monitoring and restoration activities occurred only at White Point (Figure 1). Restoration efforts began at White Point in July 2018 and are currently ongoing, as unrestored areas of the site are characterized by little to no fleshy macroalgae and high *M. franciscanus* and *S. purpuratus* densities. Reef topography is predominantly middle to low relief substrate comprised of sand, cobbles, boulders, and bedrock. The following map displays the estimated *S. purpuratus* densities before restoration activities for areas restored in Year 8 [within each 10m segment]. Site maps are also included in Appendix A.




Figure 1. Density of *S. purpuratus* (individuals per square meter) pre-restoration in White Point, Palos Verdes, California. Black square in the inset map indicates White Point location in reference to Palos Verdes. Average *S. purpuratus* density for this site is 16.66 per m², with some localized areas exceeding 150 per m² (ESRI 2021).

E) Monitoring of all Permitted sites

i. Monitoring Timeline

Table 6. Restoration and monitoring timeline July 2020 - August 2021.

| Task | 2020 | | | | | | 2021 | | | | | | | |
|------------------------|------|-----|-----|-----|-----|-----|------|-----|-----|-----|-----|------|------|-----|
| | July | Aug | Sep | Oct | Nov | Dec | Jan | Feb | Mar | Apr | May | June | July | Aug |
| Urchin Suppression | | | | | | | | | | | | | | |
| Compliance Monitoring | | | | | | | | | | | | | | |
| Response Monitoring | | | | | | | | | | | | | | |
| Analysis and Reporting | | | | | | | | | | | | | | |

Compliance Monitoring (July 2020 through June 2021)

Monitoring is conducted weekly to bi-monthly depending upon the rate of activity of restoration teams in the preceding week. Unfortunately, due to the novel COVID-19 global pandemic, active restoration of the site at White Point was disrupted. This site maintains very high *S. purpuratus* densities in the eastern portion of the cove, limiting macroalgae settlement and growth. In addition, the topography of this site consists of high relief, deep crevices, and stacked boulder complexes making restoration activities challenging. After extensive COVID-19 policy and safety protocol development, TBF biologists, staff, and commercial sea urchin harvester partners reinitiated work at White Point in September and October 2020. However, due to the location of the active restoration blocks at White Point (less than 8ft depth), combined with atypical inclement oceanic conditions (persistent wind and swell), TBF and restoration divers were often precluded from working during the remaining reporting period. TBF is adamant to take advantage of all workable weather opportunities that allow for safe and productive restoration activities. In normal circumstances, compliance monitoring work is performed by TBF biologists to ensure that restoration work is achieving performance standards. The standards are (1) the initial reduction of *S. purpuratus* to a density of two per square meter and (2) that this is being applied in a comprehensive manner sweeping through an area and not leaving patches and pockets of high *S. purpuratus* densities. All restoration areas are surveyed before and after *S. purpuratus* suppression to determine the success of restoration, and the results are entered in a georeferenced database. Post-monitoring can be completed more quickly than pre-monitoring as only the densities of *M. franciscanus* and *S. purpuratus* are counted. All 15 transects, covering 100% of the block are surveyed during post-monitoring to ensure that no pockets of high-density *M. franciscanus* and *S. purpuratus* remain at the site. Figure 2 displays the estimated *S. purpuratus* densities after restoration activities within each 10m segment of White Point. All restoration sites are re-surveyed, by roaming over the area, on a quarterly to annual basis to ensure that *S. purpuratus* densities remain at two per square meter and to observe the response of the biota to the restoration actions.




Figure 2. Density of *S. purpuratus* (individuals per square meter) post-restoration in White Point, Palos Verdes, California. Black square in the inset map indicates where the restoration area is off Palos Verdes. Average *S. purpuratus* density for this site after restoration is 1.76 per m² (ESRI 2021).

Response Monitoring (June 2020 through July 2021)

This monitoring focuses on responses of the natural community to restoration activities. The focus of this effort is subtidal utilizing an adapted Cooperative Resource Assessment of Nearshore Ecosystems (CRANE) methodology led by the Vantuna Research Group. These data provide values relating to production, species richness, and biomass.

ii. Quantity of urchins removed and collected for GSI studies and justification for removal

The estimated total number of *S. purpuratus* culled within restoration sites is 4,294,979, therefore reducing the overall average density across sites from 16.47/m² to 1.51/m². *S. purpuratus* density in some sites are less than the target density of 2/m². These low values may, in part, be attributed to habitat patchiness, physical differences among sites, and presence or accretion of fine sediment. Table 7 below shows the estimated number of urchins removed from each site by year.

Table 7. Estimated quantity of *S. purpuratus* culled by restoration site (July 1, 2013 – June 30, 2021)

Specific areas restored at Underwater Arch Cove in Years 1 and 2 were re-cleared in Years 4 and 5 due to the infiltration of *S. purpuratus* from a refuge population existing in a shallow tide pool.

| Site Name | Year 1 | Year 2 | Year 3 | Year 4 | Year 5 | Year 6 | Year 7 | Year 8 | Total by Site |
|-----------------------------|------------------|------------------|----------------|----------------|---------------|----------------|----------------|---------------|------------------|
| Underwater Arch Cove | 503,189 | 762,649 | 0 | 35,866 | 9,348 | 0 | 0 | 0 | 1,311,050 |
| Honeymoon Cove | 821,425 | 514,811 | 0 | 0 | 0 | 0 | 0 | 0 | 1,336,236 |
| Hawthorne Cove | 0 | 136,997 | 60,320 | 0 | 8,778 | 0 | 0 | 0 | 206,095 |
| Marguerite | 0 | 378,523 | 151,114 | 47,847 | 0 | 0 | 0 | 0 | 577,483 |
| Point Fermin | 0 | 0 | 160,862 | 27,263 | 6,529 | 0 | 0 | 0 | 194,654 |
| Resort Point | 0 | 0 | 0 | 0 | 49,632 | 8,559 | 0 | 0 | 58,191 |
| White Point | 0 | 0 | 0 | 0 | 0 | 330,686 | 230,479 | 50,105 | 611,270 |
| Total Urchins Culled | 1,324,613 | 1,792,979 | 372,296 | 110,975 | 74,287 | 339,245 | 230,479 | 50,105 | 4,294,979 |

Justification for Removal:

The measurement of gonad development in *M. franciscanus* and *S. purpuratus* is an important measure of secondary production in the kelp forest ecosystem and will be used to inform adaptive management of the restoration project and inform research related to kelp forests and associated fisheries. During the Year 8 reporting period, urchin collections did not occur due to volunteer restrictions put in place during COVID-19. As restrictions are lifted, TBF hopes to collect urchins for GSI studies during the Year 9 reporting period.

ii – i: Field Condition Notes

As previously mentioned, the novel COVID-19 pandemic prevented TBF biologists, staff, and our partner commercial sea urchin harvesters the ability to conduct restoration activity from April through June 2020, thus limiting any additional restoration progress at White Point. Restoration activities resumed in September 2020 and are once again on-going. As indicated elsewhere in this report and in other communication with CDFW, field conditions such as sea state, visibility, and oceanic conditions (wind and swell) were challenging for the restoration and monitoring efforts for much of 2015 and winter 2016. It is also important to note that the timing of the response monitoring for fishes and other community responses to the restoration efforts were conducted in the late spring and early summer in 2011-2014, with only two exceptions in 2011, (i.e., Honeymoon Cove and Point Vicente West were monitored on 1-28-2011 and 10-12-2011 respectively). In 2015, the surveys were conducted within the month of September except for Honeymoon Cove which was surveyed on 8-19-2015. In 2016, two rounds of surveys were conducted in spring and summer. In 2017 and 2018, all surveys were conducted in late June and July. This shift in seasonality may

affect some species differentially skewing the data. Surveys in 2019 were performed in mid-June. Surveys in 2020 were performed in late June through early August. Perhaps more significant is the strong ENSO signature elevating sea surface temperatures throughout 2015, with persistent surface temperatures off Palos Verdes neighboring 20 degrees Celsius. These abnormally high temperatures are known to affect species composition within southern California rocky reef systems. During 2020-2021, there were similar trends in several metrics between restoration and reference sites. *M. pyrifera* density, algal and invertebrate diversity, *S. purpuratus*, and fish densities/biomass were not significantly different between restoration and reference sites during Year 8 of the project. Of note however, is that *M. franciscanus* was significantly different between restoration and reference sites. Surveys revealed more observations of *M. franciscanus* in restoration sites than in reference sites; however, the abundance was still much lower than pre-restoration levels and do not indicate overpopulation.

Table 8. Response monitoring (CRANE) metadata. See Appendix B for all CRANE data tables.

| | | <u>Survey Dates</u> | | | | | | | | | | |
|--------------------|----------------------|--------------------------------|------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <u>Designation</u> | <u>Site</u> | <u>2011</u> | <u>2012</u> | <u>2013</u> | <u>2014</u> | <u>2015</u> | <u>2016</u> | <u>2017</u> | <u>2018</u> | <u>2019</u> | <u>2020</u> | <u>2021</u> |
| Restoration | Underwater Arch Cove | 2/7/11 | 6/12/12 | 6/13/13 | 7/11/14 | 9/23/15 | 6/22/16 | 7/18/17 | 6/22/18 | 6/12/19 | 6/26/20 | Aug-21 |
| | Honeymoon Cove | 1/28/11 | 3/13/12 | 5/31/13 | 7/2/14 | 8/19/15 | 6/22/16 | 7/18/17 | 6/22/18 | 6/12/19 | 6/24/20 | Aug-21 |
| | Hawthorne | 5/3/11 | 6/12/12 | 6/11/13 | 6/19/14 | 10/7/15 | 9/30/16 | 8/25/17 | 7/11/18 | 6/14/19 | 6/26/20 | Aug-21 |
| | Marguerite Central | 5/3/11 | 6/8/12 | 7/3/13 | 6/20/14 | 9/23/15 | 7/26/16 | 7/18/17 | 7/20/18 | 6/28/19 | 7/9/20 | Aug-21 |
| Reference | Ridges North | 8/12/11 | 7/17/12 | 4/26/13 | 10/29/14 | 9/11/15 | 6/3/16 | 6/30/17 | 7/11/18 | 6/12/19 | 7/23/20 | Aug-21 |
| | Rocky Point North | 6/24/11 | 6/29/12 | 7/2/13 | 7/11/14 | 9/25/15 | 6/10/16 | 6/29/17 | 7/6/18 | 6/19/19 | 7/2/20 | Aug-21 |
| | Point Vicente West | 10/12/11 | 8/10/12 | 4/24/13 | 4/18/14 | 9/23/15 | 6/22/16 | 7/25/17 | 7/18/18 | 6/14/19 | 8/14/20 | Aug-21 |
| | | <u>Bottom Temperature (°C)</u> | | | | | | | | | | |
| <u>Designation</u> | <u>Site</u> | <u>2011</u> | <u>2012</u> | <u>2013</u> | <u>2014</u> | <u>2015</u> | <u>2016</u> | <u>2017</u> | <u>2018</u> | <u>2019</u> | <u>2020</u> | <u>2021</u> |
| Restoration | Underwater Arch Cove | 15.0 | 19.0 | 15.0 | 15.8 | 21.5 | 15.0 | 18.5 | 18.0 | 15.5 | 16.0 | NA |
| | Honeymoon Cove | 15.0 | 11.5 | 18.0 | 16.5 | 18.8 | 16.2 | 20.3 | 18.3 | 15.8 | 16.0 | NA |
| | Hawthorne | 14.4 | 19.0 | 17.0 | 17.0 | 21.0 | 18.0 | 16.8 | 20.6 | 16.0 | 15.0 | NA |
| | Marguerite Central | 15.0 | 17.0 | 17.0 | 20.0 | 22.0 | 14.0 | 20.0 | 20.0 | 18.0 | 19.7 | NA |
| Reference | Ridges North | 18.0 | 16.6 | 13.7 | 19.8 | 21.0 | 15.0 | 17.9 | 22.0 | 16.5 | 12.6 | NA |
| | Rocky Point North | 18.0 | 15.0 | 18.0 | 21.0 | 21.0 | 14.3 | 16.8 | 19.5 | 16.5 | 17.0 | NA |
| | Point Vicente West | 11.0 | 19.0 | 13.2 | 13.5 | 21.0 | 15.2 | 19.7 | 19.5 | 16.5 | 16.2 | NA |
| | | <u>Coordinates</u> | | | | | | | | | | |
| <u>Designation</u> | <u>Site</u> | <u>Latitude</u> | <u>Longitude</u> | | | | | | | | | |
| Restoration | Underwater Arch Cove | 33.75291 | -118.41499 | | | | | | | | | |
| | Honeymoon Cove | 33.76459 | -118.42406 | | | | | | | | | |
| | Hawthorne | 33.75068 | -118.41558 | | | | | | | | | |
| | Marguerite Central | 33.75694 | -118.41772 | | | | | | | | | |
| Reference | Ridges North | 33.78697 | -118.42065 | | | | | | | | | |
| | Rocky Point North | 33.77966 | -118.42739 | | | | | | | | | |
| | Point Vicente West | 33.74073 | -118.41283 | | | | | | | | | |

iii. Species Richness

Species richness is the number of unique species found at a site. The species richness values are derived from the CRANE surveys provided by VRG. Since restoration events, species richness has increased in all restored sites (Table 9). Though these values are slightly variable from year to year, the restored sites post 2013 (post 2015 for Marguerite Central) do have similar richness values when compared to reference sites.

Table 9. Fish Species Richness (total number of species).

| Site Type | Site | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 |
|----------------|----------------------|------|------|------|------|------|------|------|------|------|------|------|
| Kelp Reference | Point Vicente West | 8 | 6 | 10 | 11 | 12 | 13 | 9 | 11 | 10 | 11 | 8 |
| | Ridges North | 6 | 11 | 7 | 6 | 5 | 10 | 5 | 12 | 8 | 7 | 5 |
| | Rocky Point North | 8 | 8 | 8 | 8 | 6 | 7 | 9 | 11 | 8 | 4 | 6 |
| Restoration | Hawthorne | 10 | 6 | 8 | 7 | 10 | 13 | 11 | 12 | 12 | 7 | 9 |
| | Marguerite Central | 6 | 10 | 10 | 9 | 10 | 10 | 7 | 9 | 11 | 9 | 11 |
| | Underwater Arch Cove | 6 | 9 | 6 | 12 | 7 | 8 | 11 | 9 | 9 | 8 | 8 |
| | Honeymoon Cove | NA | 2 | 4 | 8 | 5 | 12 | 7 | 8 | 8 | 5 | 5 |

iv. Density of Kelp Forest and Ecosystem Species

As a measure of kelp forest density, we analyze the number of stipes per 100 m² that are greater than one meter in height. The *M. pyrifera* stipe density is provided by VRG during their annual CRANE surveys. The years after post restoration activities (2016-2021) showed an immediate increase in the *M. pyrifera* stipe density for all four restoration sites (Figure 3). Increases in stipe density post-2015 are orders of magnitudes higher than the years prior to restoration (2011-2014). *M. pyrifera* densities increased in 2020 across both restoration and kelp reference sites. *M. pyrifera* densities decreased in 2021 across both restoration and kelp reference sites, with the exception of the Hawthorne restoration site, which showed increases in density compared to 2019 and 2020. Differences in stipe density post-restoration are likely explained by natural inter and intra-annual variation; e.g., kelp canopy cover, transmissivity, temperature, nutrient availability, and upwelling. It should be noted that restoration events did coincide with a natural mass mortality event that contributed to decreased urchin density.




Figure 3. *Macrocyctis pyrifera* stipe density (individuals per 100 m²). Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *M. pyrifera* was not significantly different by site designation in 2021 ($t= 1.33$, $p=0.24$).

Kelp Canopy Area and Percent Cover by Site

Since 2003, MBC Aquatic Sciences has been hired by the Central Region and Region Nine Kelp Survey Consortium to take quarterly aerial surveys of the mainland Southern Californian kelp forests. These kelp surveys inform the consortiums about the status of the kelp forests and serve to determine possible impacts that dischargers and environmental variables are having on the kelp beds. These surveys consist of digital color and infrared color photos taken of the kelp beds that are then processed into base maps. These surveys cover approximately 354 km of the 435 km southern Californian coastline from Ventura to the U.S./Mexico boarder (MBC 2018).

The consortiums provided TBF with the base maps of annual kelp bed maximums of the Palos Verdes kelp beds, which can be used to show the progress of restoration off Palos Verdes. Surveys from 2011 through 2015 show an overall increase in kelp canopy acreage off the peninsula; however, kelp canopy dropped in 2016 due to the ENSO event. In 2017, sea conditions returned to more normal state and kelp canopy started to recover in areas where kelp was eliminated during the ENSO event. In 2018, MBC reported kelp beds off Palos Verdes increased substantially, some of which grew to levels approaching the maximum acreage observed since surveying began (MBC 2018).

At the close of 2019, MBC, the Central Region and Region Nine Kelp Survey Consortium made the decision to change the reporting frequency from annually to biennially. Therefore, the data typically made available to TBF in past years was unavailable for 2019. Thus, in the Year 7 report, no overlay of the MBC data canopy cover with TBF restoration sites was possible. MBC and the Kelp Survey Consortium only released the report a few weeks prior to the due date of this report. TBF did not have sufficient time to process, analyze and generate the maps previously generated for this report.

M. franciscanus* and *S. purpuratus

Both *M. franciscanus* and *S. purpuratus* densities began declining in 2013-2014 (Figures 4 & 5). Their numbers remained low until the end of Year 5 when pulses of urchins (mainly *S. purpuratus*) were observed in several areas off the peninsula. Although CRANE surveys show a sharp decline prior to restoration activities at Marguerite Central, TBF fine-scale density data shows that our restoration efforts did decrease purple urchin high-density patches further between 2014-2016. Decreases prior to restoration activities could possibly be a result of early effects of the observed 2014-2015 natural wasting event, or discrepancies in CRANE surveying. TBF suspended *S. purpuratus* suppression from the fall of 2015 through the spring of 2016 to monitor the wasting event. Suppression continued in the late spring of 2016 once lesions on *M. franciscanus* and *S. purpuratus* were no longer found and densities of greater than 2/m² persisted within our restoration sites. *M. franciscanus* densities also dropped during this time, even though TBF does not suppress this species. The decline in abundance was most likely caused by three factors, (1) *M. franciscanus* and *S. purpuratus* wasting event, (2) commercial sea urchin harvesters extracting the *M. franciscanus* for the fishery, and (3) an increase in cryptic behavior. A small uptick in *M. franciscanus* and *S. purpuratus* were recorded during community analysis surveys for Year 5 for both species. In Year 6, we saw a small increase in *M. franciscanus* density and a small decrease in *S. purpuratus* density. In Year 7, we saw a small decrease in both *M. franciscanus* and *S. purpuratus* densities. In Year 8, at restoration sites we saw increases in the number of *S. purpuratus* observed in Hawthorne and Underwater Arch Cove, but negligible change at Honeymoon Cove and Marguerite sites. At reference sites, slight decrease was observed at Point Vicente West, while slight increases were observed at Ridges North and Rocky Point North sites. As for *M. franciscanus* in Year 8, *M. franciscanus* was significantly different between restoration and reference sites, with restoration sites showing slight increases, while reference sites showing declines or negligible change. Surveys revealed an increase in

observations of *M. franciscanus* in all restoration sites, however the abundance was still much lower than pre-restoration levels and do not indicate overpopulation.




Figure 4. *M. franciscanus* density (individuals per 100 m²). Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *M. franciscanus* density was significantly different by site designation in 2021 ($t= 3.24$, $p= 0.045$).




Figure 5. *S. purpuratus* density (individuals/100 m²). Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *S. purpuratus* density was not significantly different by site designation in 2021 ($t= 1.77$, $p = 0.175$).

Panulirus interruptus

Panulirus interruptus (California Spiny Lobster) were quantified in CRANE invertebrate swaths. Prior to *S. purpuratus* removal in restoration sites, *P. interruptus* were not found within the sites (Figure 6). There has been a notable increase in the abundance of *P. interruptus* within restoration sites starting in 2016. While the abundance in restoration sites declined in 2019, the population observed remains larger than pre-restoration abundance levels. In 2020, the population in restored areas exceeded the population observed in reference sites. In 2021, the population in reference sites was slightly larger than the population observed in restored sites. It should be noted, however, *P. interruptus* abundance is highly variable among sites and years, exemplified by the decline in population in kelp forest reference sites as well. This decline could be attributed to two factors: (1) commercial lobster fishing pressure is heavy throughout the Palos Verdes region, (2) *P. interruptus* are mobile and can select for areas based off preferable habitat and oceanographic conditions.




Figure 6. Mean *P. interruptus* density (Individuals per 100 m²) at kelp forest reference sites shown in green (Ridges North, Rocky Point North, and Point Vicente West), and restoration sites shown in blue (Underwater Arch, Honeymoon Cove, Marguerite Central and Hawthorne).

v. Density and biomass of kelp bass and California sheephead

Fish Data Processing

Sites were sampled over a period of several months and seasons, thus, young-of-the-year (YOY) were removed prior to fish density calculations because they could numerically dominate the assemblage at some sites sampled early in the season, but decline later in the year due to natural mortality. YOY were generally defined as fishes <10 cm, except for some smaller species, where they were defined as individuals less than between 1.5 and 5 cm based on published species-specific growth rates and expert opinion. Total length (TL) estimates were converted to biomass using standard species-specific length-weight conversions from the literature. YOY were not excluded from biomass calculations, as their small size will influence biomass estimation less than abundance estimation. Density and biomass was then summed across all three portions (bottom, midwater and canopy) of each transect, except for when the water depth is less than 6m, meaning that the volumes of the canopy and midwater portions would overlap, in which case no midwater portion was included. Density values were then scaled to the number per 100m².

Paralabrax clathratus (kelp bass) abundance and biomass has gradually increased in restoration sites since restoration efforts were started (Figures 7 & 9). In the surveys conducted in 2018, kelp bass density and biomass indicate an increasing trend since being restored and are on par with kelp forest reference sites. This increased number of kelp bass could be due to a multiyear increase and persistence of *M. pyrifera* within these restoration sites. During 2020 surveys, the overall kelp bass density for reference sites increased, while there was a slight decline in overall density for restoration sites. This is punctuated by a steeper decline from Marguerite Central from 2019 to 2020, most likely associated with impacts from the increased sedimentation on the reef resulting from the coastal bluff slough that occurred in spring 2019. During 2021 surveys, *P. clathratus* increased density at Underwater Arch, Marguerite, Hawthorne and Ridges North Sites, while Honeymoon, Rocky Point North, and Point Vicente West decreased. Trends for biomass were similar to density yet, Underwater Arch showed a decline, while Rocky Point North showed an increase. Overall, density and biomass in restoration sites depict the same trends as kelp reference sites, with no significant difference by site type.

Kelp bass recruit to kelp canopy and use kelp as a refuge to hide from predators or to ambush prey. Biomass of kelp bass from 2020 shows that the largest biomass of kelp bass is within Point Vicente MPA site, which is markedly higher than other reference and restoration sites. This is expected as fishing is not allowed within this area, allowing for fish to grow larger without fishing pressure. All current restoration sites are outside MPAs where fishing is allowed. Restoration sites may have a larger density compared to reference sites, yet smaller biomass, on account of fishing pressure for larger sized individuals, thus leaving a high abundance of smaller sized fish in restoration sites. In 2021, kelp bass biomass at Point Vicente declined dramatically, leaving 2021 results indicating that the restoration sites show similar, if not elevated, biomass values compared to reference sites.

Semicossyphus pulcher (California sheephead) abundance and biomass has been variable among monitoring years for all sites (Figures 8 & 10). Surveys from 2021 continue to exhibit this annual variation, with Underwater Arch, Honeymoon, Marguerite, Rocky Point North, and Point Vicente West sites displaying slight increases in density and biomass from 2020, while Hawthorne and Ridges North sites show slight decreases. However, density and biomass in restoration sites depict the same trends as kelp reference sites, with no significant difference by site type.




Figure 7. Density of *P. clathratus* by site type: restoration and reference. Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *P. clathratus* density was not significantly different by site designation in 2021 ($t = 2.41, p = 0.083$).




Figure 8. Density of *S. pulcher* by site type: restoration, and reference. Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *S. pulcher* density was not significantly different by site designation in 2021 ($t = -0.04$, $p = 0.973$).




Figure 9. Biomass of *P. clathratus*, per 100 m², by site type: restoration, and reference. Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *P. clathratus* biomass was not significantly different by site designation in 2021 ($t = 1.7$, $p = 0.167$).




Figure 10. Biomass of *S. pulcher*, per 100 m², by site type: restoration, and reference. Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. *S. pulcher* biomass was not significantly different by site designation in 2021 ($t = 0.12$, $p = 0.914$).

Community Diversity

The Shannon-Wiener diversity index came from information theory and measures the order (or disorder) observed within a particular system. The Simpson's index of diversity accounts for both richness and proportion of each species. It has been a useful tool to terrestrial and aquatic ecologists. Both diversity measures show a rapid increase of algal/invertebrate diversity once restoration was completed in Underwater Arch, Honeymoon Cove, and Hawthorne (Figure 11). After restoration activity, diversity measures show little fluctuation, apart from Marguerite Central, as it appears diversity decreased slightly in the year after restoration was completed. In 2021, restoration sites mimic diversity index measurements for kelp reference sites for both algal/invertebrate diversity and fish diversity (Figure 11 & 12).




Figure 11. Algal and invertebrate diversity at Restoration sites (Underwater Arch, Honeymoon Cove, Marguerite Central and Hawthorne) and Reference sites (Ridges North, Rocky Point North, and Point Vicente West). Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. Diversity measures used are Shannon-Wiener ($t= 2.08$, $p= 0.098$) (Left) and Simpson's Diversity ($t= 1.57$, $p= 0.196$) (Right).




Figure 12. Fish diversity at Restoration sites (Underwater Arch, Honeymoon Cove, Marguerite Central and Hawthorne) and Reference sites (Ridges North, Rocky Point North, and Point Vicente West). Sites Underwater Arch, Honeymoon Cove, and the majority of Hawthorne were restored as of 2015. Restoration began in 2015 at the site Marguerite Central (previously a control site) and was completed in the Spring of 2017. Diversity measures used are Shannon-Wiener ($t= 0.02$, $p= 0.988$) (Left) and Simpson's Diversity ($t= 0.18$, $p= 0.863$) (Right).

vi. Gonadosomatic indices of *M. franciscanus* and *S. purpuratus*

The measurement of gonad development in *M. franciscanus* and *S. purpuratus* is an important indicator of secondary production in the kelp forest ecosystem and is used to inform adaptive management of the restoration project and research related to kelp forests and associated fisheries. The gonadosomatic index is the ratio of the weight of the gonad to the overall weight of the animal.

No urchins were collected for the Year 8 annual report. In order to process urchins in a timely manner (to reduce stress and water loss from their gonads), collection and dissection requires a large effort consisting of student and community volunteers. In previous years, TBF divers were able to collect urchins at one kelp reference, two restoration, and one barren control site before transporting all urchins to LMU. More than 50 student and community volunteers would then process urchins throughout the day/night. Due to COVID-19 restrictions with organizing large groups of people, as well as LMU closing lab spaces, TBF was not able to hold this event in 2020.

- F) Analysis of the ecosystem response to the restoration activities at the restoration sites, including species that are key indicators of a healthy and persistent kelp forest ecosystem.

Community Analysis Methods

As part of the quantitative characterization of the community structure of the reefs, we examined patterns in the overall kelp forest community using fish and swath (benthic macroinvertebrates and kelps) data combined. Density metrics were square root transformed (fish and swath data). Two-dimensional, non-metric multidimensional scaling (nMDS) was used to examine patterns among kelp forest communities (Figure 18) and fish biomass (Figure 19) at sites using the 'metaMDS' function in the 'vegan' package (Oksanen et al. 2016) in R (R Core Team 2016). A similarity matrix constructed with transformed taxon-specific values (site means for each site/sampling period combination) and the Bray-Curtis similarity. To provide context to the observed relationships amongst sites, patterns of taxa densities were visualized across the nMDS ordination plots using the 'ordisurf' function in the R package 'vegan' (Oksanen et al. 2016) which fits a smooth surface using generalized additive modeling (GAM) with thin plate splines (Wood 2003, Oksanen et al. 2016). These visualizations help inform drivers of community structure as seen in nMDS plots.




Figure 18. (Left) Two-dimensional Non-metric multidimensional ordination plot of kelp and macroinvertebrate communities using Bray-Curtis similarity based on the square-root transformed mean taxa density for each site/sampling period combination. (Right) Two-dimensional nMDS plot of fish density for each site/sampling period combination. Open circles indicate every site sampled, while closed circles indicate the mean values for the site type.




Figure 19. Two-dimensional Non-metric multidimensional ordination plot of fish biomass using Bray-Curtis similarity based on the square-root transformed mean taxa density for each site/sampling period combination. Open circles indicate every site sampled, while closed dots indicate the mean values for the site type. Fish communities depict an evolution of restoration sites, forming a large significant cluster near kelp reference sites, which are visibly differentiated from pre-restoration values.

Community Analysis Results

The three plots presented above display a convergence over time in which restoration sites begin to resemble, structurally, the reference sites after purple urchin density reduction. The earlier years depicted in these plots show that the converse was true in advance of restoration efforts; that the structure of restoration sites, pre restoration, resembled control sites (sites that contained urchin barrens for comparison early in the project).

Two restoration sites were completed near the close of 2014. The community analyses show a convergence of restoration and reference sites in 2014 as the restoration sites changed from barrens to young kelp forests. The occurrence of a mass wasting event of *M. franciscanus* and *S. purpuratus* happened with considerable severity off the Palos Verdes Peninsula impacting reference and restoration sites in 2015 into 2016. This further loss of top down pressure from *M. franciscanus* and *S. purpuratus* on the development of *M. pyrifera* and other macroalgae and the freeing from competition, of other grazers, likely caused this progression from barren to young kelp forest to continue in 2015-2016.

These plots indicate, with confidence, that the loss of *M. franciscanus* and *S. purpuratus* i.e. a reduction in their density, allows for the growth and development of other benthic organisms that are no longer limited by the direct and indirect impacts of *M. franciscanus* and *S. purpuratus* grazing. Further

monitoring of these sites may, over time, detect trends that elucidate more subtle or developing relationships in community structure. Likely, these characteristics will be displayed via divergence of these site types over time, or in response to other forms of disturbance and other stressors.

The plots also support the idea that *S. purpuratus* suppression creates similar near-term changes in community structure to widespread reductions in *M. franciscanus* and *S. purpuratus* due to disease. These different causes of *M. franciscanus* and *S. purpuratus* density reduction have both driven formerly barren reef states to resemble reference sites (i.e. sites with persistent kelp and more complex community structure). These results suggest that in the near-term, *S. purpuratus* suppression is a fair mimic for natural losses in *M. franciscanus* and *S. purpuratus* populations driving kelp forest community structure on a local scale. See Williams et al. 2021 for further discussion on the community analysis depicting the convergence of structural metrics of restoration sites to resemble reference sites.

G) Evaluation of successes and failures of restoration activities for each site

A few statements can be made that generally describe conditions during this project that directly impacted the amount, type, and accuracy of work conducted. 2015-2016 proved to be one of the most powerful El Niño signatures recorded on the west coast of the United States. This El Niño event followed, and was perhaps strengthened, by the persistence of “the blob”, a large area of atypically warm ocean surface water that impacted the California Current. For Palos Verdes and elsewhere in southern California, these environmental factors resulted in abnormally high sea surface temperatures, which were only punctuated periodically by localized upwelling events. The thermal related stress associated with the confluence of these stressors slowed or prevented the development of *M. pyrifera* and other macroalgae and may have contributed to the virulence and mass wasting of several genera of *Pisaster* spp. and in the fall of 2015, a seemingly similar, yet less widespread or virulent wasting of *M. franciscanus* and *S. purpuratus*. In 2016, the project failed to collect a sufficient number of *M. franciscanus*, and individuals were not collected from barren sites for dissections. During 2019, the project was only able to collect one *M. franciscanus* from a barren site. There are currently no signs of widespread mass wasting disease off Palos Verdes. During pre and post monitoring surveys, divers utilize flashlights to more accurately and efficiently quantify *M. franciscanus* and *S. purpuratus* in active restoration sites.

During 2017-2018 (Year 5 of the project), *M. franciscanus* and *S. purpuratus* densities rose noticeably, and TBF chose several areas of concern to begin restoration efforts. These sites (mainly Resort Point and a new area of Hawthorne, as well as minor work in Underwater Arch Cove and Point Fermin) consisted of high *S. purpuratus* densities, but also supported high biomass of fish, invertebrates, and *M. pyrifera*. The work done in these sites were to cull *S. purpuratus* numbers to prevent these reefs from reverting to pre-2015 barrens. Near the beginning of summer 2018 more reef was identified as having high *S. purpuratus* densities, and White Point was chosen as the most problematic. This site had developed into an *S. purpuratus* barren, devoid of macroalgae besides coralline algae, and a 2018 average density of 24.24 *S. purpuratus* per m², with some isolated patches reaching densities of 150+ per m². TBF began monitoring and restoring this site in the summer of 2018. Restoration at White Point is in progress but incomplete as of the close of the Year 8 timeframe. Inclement weather conditions, in conjunction with COVID-19 restrictions, precluded significant restoration activities to occur in Year 8. Further progress is underway and planned for Year 9 reporting period.

Underwater Arch Cove

Underwater Arch Cove was considered restored in January 2015, being that no expanses of the reef were observed to support densities of *S. purpuratus* in excess of two per square meter. However, in Year 4 of the project, one locale within Underwater Arch Cove showed higher than two *S. purpuratus* per square meter during the spring and summer of 2016. The Bay Foundation re-monitored Underwater Arch to determine the expansion of *S. purpuratus* in the area and to decide if suppression should be started again. The expansion of *S. purpuratus* was found to be relatively contained near the large tidepool at the north edge of the site. The renewed restoration of this section of Underwater Arch took place from 4/7/17 – 6/20/17, which reduced *S. purpuratus* densities from 4.83/m² to 1.07/m² across a total area of 2.34 acres.

In Year 5 of the project, the Underwater Arch restoration site was similar to kelp reference sites in terms of *M. pyrifera*, fish biomass, and *S. purpuratus* densities. However, during annual monitoring at the end of Year 5, increased densities were observed further east of the area revisited in Year 4. TBF biologists culled *S. purpuratus* from 0.28 acres on 7/6/17. This area was surveyed in the fall of 2018 and may require *S. purpuratus* suppression to reestablish the kelp forest. At this time, no additional restoration efforts have been conducted at the site, but divers will continue to monitor the area and assess *S. purpuratus* densities. In Year 8, we observed small pockets of reef exhibiting high *S. purpuratus* densities. Analyses showed fish community diversity (both Shannon-Wiener and Simpson's indexes) decreased. In addition, algae/invertebrate community diversity (both Shannon-Wiener and Simpson's indexes) decreased from Year 7. However, in both cases, the site maintained values well above pre-restoration. We plan to closely observe urchin density in Year 9 in the case that restoration actions be taken.

In July 2021, georeferenced photos and video were taken to visually represent the changes over time at Underwater Arch.

Honeymoon Cove

Honeymoon cove restoration was completed in January 2015. Surveys have since been conducted by TBF personnel periodically to quickly assess the condition of the reefs found in this cove. This will continue in the coming year to ensure that the restoration target of two *S. purpuratus* per square meter are maintained and that *M. pyrifera* and other biota are persisting in the area. Due to the high success of restoration at Honeymoon Cove, The Bay Foundation and NOAA biologists outplanted 827 *Haliotis fulgens* (Green Abalone) onto a section of restored reef in June 2015. Subsequent monitoring was conducted in March of 2017 identifying several emergent *H. fulgens* on the site. The last survey completed in August 2021 found approximately 89 *H. fulgens* within the 10 by 10-meter outplant site. Prior to outplanting, only 10 *H. fulgens* were found within the site. Genetic analysis based upon tissue samples taken in situ is ongoing to determine whether these emergent *H. fulgens* are in whole or in part the same organisms that were outplanted.

The increase in biomass of *P. clathratus* has been gradual since restoration completion. Fish and algae/invertebrate community diversities (both Shannon-Wiener and Simpson's indexes) have been variable year to year, but the general trend depicts increases since restoration, with year 8 displaying higher values than year 7.

Georeferenced photos and video were collected in August 2021 to document conditions within the site over time. See Appendix C.

Resort Point

Resort Point is a deep site (40-60 feet) located offshore of Honeymoon Cove. Resort Point has had persistent kelp since the start of this project, yet consisted of high *M. franciscanus* and *S. purpuratus* densities. The reef complex that exists between Honeymoon Cove and Resort Point has no barrier to *M. franciscanus* and *S. purpuratus* movement e.g., sand channels and high energy shallow prominences. Restoration blocks along the western edge of Honeymoon Cove were separated by a thin line of existing kelp approximately 45 m wide. Both the continuity of this high concentration of urchins to Honeymoon Cove and lack of an incursion barrier encouraged urchin suppression at Resort Point. Urchin suppression focused on protecting this existing kelp forest in a total of 4 acres. Divers will continue to visit this site periodically to ensure densities have not increased.

The greater average depth of Resort Point and the presence of kelp make it a statistical outlier in the overall monitoring scheme developed to inform the project. Thus, to determine the effect of the work conducted at Resort Point and Honeymoon Cove, we have relied on the data collected within Honeymoon Cove, consistent with other depth profiles, for comparability across restoration and reference sites. Though separately defined operationally, we consider these two sites to function statistically as a single unit.

Photo points and video transects were not established in this site due to its proximity to Honeymoon Cove.

Hawthorne

The Hawthorne restoration site exists south of Underwater Arch Cove. It is a section of exposed coast comprised of large bedrock shelves, as well as boulders forming low lying expanses of unconsolidated reef. In spring 2017, an area slightly less than an acre (0.89 acres) outside of the previously restored area was found supporting high densities of *S. purpuratus* and was cleared during the summer of 2017. Hawthorne has proven to be a very dynamic site with high wave energy and considerable sediment movement. Consequently, the neighboring reef that initially surrounded the large rock and pinnacle in the HAW 2 block (where the permanent photo point is located), has been covered as a result of sediment transport.

In Year 8, fish community diversity (both Shannon-Wiener and Simpson's indexes) displayed a decrease, while algae/invertebrate community diversity (both Shannon-Wiener and Simpson's indexes) increased slightly, maintaining values significantly higher than pre-restoration values.

Georeferenced photos and video were collected in July 2021. See Appendix C.

Marguerite

At the start of this project, Marguerite was designated to serve as a control (barren) site throughout the permitted work. However, in 2015, discussions with CDFW resulted in the expansion of restoration actions to Marguerite. Marguerite is an expansive area of reef located between Honeymoon Cove to the north and Underwater Arch Cove to the south. Restoration actions were initiated in December of 2015 at the southern and northern terminuses of this site. At times, three restoration teams were working in this area as they progressed towards one another reducing the gap between them. This site is openly exposed to northerly and westerly swell energy and receives some wrap around from south westerly energy. This site is comprised of high relief reef with semi vertical walls, 20-30 feet in height extending from the sea floor to the surface. Between these reefs, expansive boulder fields and some sandy expanses exist. The shoreline is defined by bench-like bedrock or cobble beaches. The physical structure

of this site supports higher rates of fish production and increased diversity of benthic organisms due to its heterogeneity.

Although Figures 4 and 5 show a complete lack of *M. franciscanus* and *S. purpuratus* throughout the 3 years of restoration activities (2015-2017), the fine-scale density surveys completed by TBF in the Marguerite site showed that densities of purple urchins remained above 2/m². Year 2 pre-restoration density of *S. purpuratus* was 19.52/m², Year 3 pre-restoration density of *S. purpuratus* was 11.06/m², and Year 4 pre-restoration density of *S. purpuratus* was 3.33/m², all representative densities associated with urchin barrens. In Year 8, fish community diversity (both Shannon-Wiener and Simpson's indexes) displayed an increase, while algae/invertebrate community diversity (both Shannon-Wiener and Simpson's indexes) decreased. Fish diversity has remained high at this site throughout monitoring due to its physical structure. Algae/invertebrate diversity remains significantly above pre-restoration values.

The photo and videos for Marguerite were collected in July 2021. However, of note, extremely poor visibility inhibited video quality in the inner portion of the site. Visibility was hampered by sediment on the reef.

Point Fermin

Point Fermin is near the southeastern terminus of the Palos Verdes Peninsula. Restoration actions were started in July of 2015 and continued through February 2016 but were then suspended until October 2016. Approximately four (3.93) acres were restored in those few months, clearing a barren that is roughly central to the shallow expanses of the reef complex. Restoration activities resumed in October 2016 through December 2016, clearing 1.35 acres of reef. An additional 0.22 acres were restored in July 2017. This area is low relief, largely tabular, and dominated by soft sedimentary rock, making it very different from the other restoration sites. In addition, Point Fermin is sheltered from northerly and westerly swells, but is directly exposed to southerly and some south westerly wave energy. The restoration efforts at Fermin have resulted in the development of *M. pyrifera*, other macroalgae and several phyla of sessile life on the reef. While initial restoration activities yielded positive results, culminating with a 98% canopy cover in 2017 according to MBC data, during the Year 6 reporting period, TBF staff discovered high densities of *S. purpuratus*, resulting in a shift back towards a barren state. This phase shift can best be explained by two factors: (1) during restoration efforts a large *S. purpuratus* recruitment pulse occurred where many individuals were observed in the 0.5-1cm range making comprehensive and targeted suppression difficult, and (2) the site contains several long channels inshore with deep crevices occupied with larger *S. purpuratus*. Therefore, TBF staff speculate that intrusion from this area into the site may have contributed to the shift back towards a barren state.

Similar to White Point, CRANE surveys were never conducted for Point Fermin, due to different exposure, substrate characteristics, and the unsettled condition of the site from start and stop restoration actions. As restoration work is required to address the current barren state of the reef at Point Fermin and proposed to continue for Year 9 of the project, CRANE surveys will be conducted moving forward, beginning in 2022. Data collected and analyzed for the restoration efforts at Point Fermin will be used solely to describe the condition and trends within this site over time.

Photos and videos for Point Fermin were collected in June 2021. The photos and videos from Point Fermin, in previous years, quite convincingly display the changes resulting from the *S. purpuratus* suppression in that site, and further display the consequences of refuge *S. purpuratus* populations. Additional area at Point Fermin will be targeted during Year 9 of the project.

White Point

White Point is a section of exposed coast located north of Point Fermin and has a depth profile that ranges from 5-35 feet. The White Point restoration site was established in the summer of 2018 due to the high density of *M. franciscanus* and *S. purpuratus*. At the close of the Year 6 reporting period, 3.11 acres out of an estimated 9.93 acres have been cleared of excess *S. purpuratus*. In Year 7 of the project, an additional 4.38 acres were restored, and further surveys expanded the estimated barren area from 9.93 to 15 acres. The substrate of the area is primarily comprised of bedrock and scattered boulder cover, with various pinnacles becoming exposed during low tide. In addition, there are significant sand channels interspersed within the site. Community response monitoring for White Point was delayed a year due to funding and was monitored for the first time in August 2020.

CRANE surveys were never conducted within White Point due to the proximity of standing kelp along the perimeter of the barren and the exposure of the site. The distance identified at the outset of the project for the entirety of two CRANE transects, at fixed depths, comprising 60 meters in length, was to be at least 20 meters from the edge of existing kelp. The configuration of the barren at White Point, and the topography, did not accommodate the necessary distance and would have skewed the results. In 2020, CRANE surveys were conducted for the first time as the distance to existing kelp was determined to be sufficiently distant to not compromise the data, due to edge effects. In addition, White Point and Point Fermin share a different exposure than the near contiguous efforts located further to the north and west. This difference may cause localized responses due to differences in wave energy, upwelling, wind, turbidity, and (based upon observations) transmissivity and possible recruitment events. For these additional factors, data will be collected moving forward but will not be analyzed to determine the overall trend of restoration sites at Palos Verdes. The data will solely be used to describe the condition of the restoration site at White Point over time. As evident in Table 10 below, increases in fish density and biomass, swath diversity, and giant kelp density were apparent from 2020 to 2021, showing the beneficial aspects of restoration activities at this site.

Table 10. Community analysis monitoring data for White Point.

| White Point Crane Data - September 2020-2021 | | |
|--|---------------|---------------|
| Analysis | 2020 | 2021 |
| Coordinates: | | |
| Latitude | 33.71287 | 33.71287 |
| Longitude | -118.3159 | -118.3159 |
| Temperature (°C) | 17.5 | 16.1 |
| Fish Richness | 10 | 10 |
| Fish Diversity H | 2.029 | 0.48 |
| Fish Diversity 1-D | 0.83 | 0.17 |
| Fish Density: | | |
| <i>Paralabrax clathratus</i> (/100m ²) | 2.1 ± 0.4 | 3.3 ± 2.3 |
| <i>Semicossyphus pulcher</i> (/100m ²) | 1.3 ± 0.4 | 1.3 ± 0.4 |
| Fish Biomass: | | |
| <i>Paralabrax clathratus</i> (g/100m ²) | 373.3 ± 108.5 | 514.3 ± 350 |
| <i>Semicossyphus pulcher</i> (g/100m ²) | 402.2 ± 220.6 | 572.4 ± 572.4 |
| Swath Diversity H | 0.875 | 1.997 |
| Swath Diversity 1-D | 0.35 | 0.832 |
| Swath Density: | | |
| <i>Macrocystis pyrifera</i> stipes (/100m ²) | 11.7 ± 8 | 687.5 ± 60.8 |
| <i>Panulirus interruptus</i> | - | 1.67 ± 1.67 |
| <i>Mesocentrotus franciscanus</i> (/100m ²) | 4.2 ± 4.2 | 15 ± 3.3 |
| <i>Strongylocentrotus purpuratus</i> (/100m ²) | 510 ± 278 | 160.9 ± 6.3 |

Photos for White Point were collected in June 2021, depicting the establishment of understory algae (*Dictyopoteris* sp., *Zonaria* sp., *Eisenia arborea*, and other brown and red algae), as well as canopy *M. pyrifera*, indicating the successful transition away from a barren state. The White Point permanent video transect was established in summer 2019. Restoration work will continue into Year 9.

Note: Figure 20 below displays all the restoration sites on the same map for a comprehensive look at the scale of the project in relation to the Palos Verdes peninsula.




Figure 20. Urchin barrens as mapped in 2010 and areas restored, representing a possible expansion and/or shift of urchin barrens. The locations of urchin barren areas are in pink, restoration areas completed in Years 1 through 7 are green, and restoration areas in Year 8 are blue (ESRI 2021).

H) Geo-referenced images before and after restoration activities

Between July 1, 2013 and August 31, 2021, photos and video were taken at various locations within six restoration sites both pre and post restoration efforts (excludes Resort Point extension of HMC site). The GPS coordinates and maps displaying locations of these photos and videos are listed in Appendix C.

Permanent photo points have been identified in six sites, which will be photographed over time. These locations were chosen due to either, a unique geological feature, or frequency of diving due to other projects occurring in the area. Some sites have distinct, recognizable rock structures, but once kelp recruits back into the area these features are often obscured. Video transects were also established in each site starting from a known GPS coordinate and laying 30m transect tapes at a predetermined heading. The paths of these video transects and photo points are mapped in Appendix C. We aim to increase our efficiency by revisiting the permanent photo points and a select subset of transects for video at minimum once per year during late summer to early winter (June to November), providing an overview of the conditions and response within each site. Full video transects for 2021 have been recorded and time-lapse videos were edited together using one 30m segment to show changes over time within each site.


Table 11. Permanent photo point selections in restoration sites.

| Restoration Site | Latitude | Longitude | Notes |
|------------------------------|----------|-----------|--|
| Honeymoon Cove - T2 | 33.76426 | -118.4237 | East-west running ridge |
| Honeymoon Cove - R5 | 33.7653 | -118.4242 | <i>Haliotis fulgens</i> outplant site monitored annually |
| Marguerite - T16 | 33.75756 | -118.4178 | Annual surveys conducted |
| Underwater Arch J1 - J2 - T7 | 33.7526 | -118.4146 | Original video transect, repeated annually |
| Hawthorne - T2 | 33.75064 | -118.4161 | Large pinnacle within block 2 |
| Point Fermin - J7 | 33.70303 | -118.2902 | North-south running ridge |
| White Point - T12 | 33.71297 | -118.3165 | Large boulder 7meters 0 degrees from block 12 smile |

Literature Cited


- ESRI. 2021. ArcGIS Desktop: Release 10.8.1. Environmental Systems Research Institute, Redlands, CA.
- Ford T, Meux B. 2010. Giant Kelp Community Restoration in Santa Monica Bay. *Urban Coast 2*: 43-46.
- MBC Aquatic Sciences. 2018. Size of the Kelp Beds in 2018: Ventura, Los Angeles, Orange & San Diego counties. Central Region Kelp Survey Consortium and Region Nine Kelp Survey Consortium. Costa Mesa, CA: MBC Aquatic Sciences.
- Williams, J.P., Claisse, J.T., Pondella II, D.J., Williams, C.M., Robart, M.J., Scholz, Z., Jaco, E.M., Ford, T., Burdick, H. and Witting, D., 2021. Sea urchin mass mortality rapidly restores kelp forest communities. *Marine Ecology Progress Series*, 664, pp.117-131.

Appendix A: Palos Verdes Kelp Restoration Project Map Images




Map A1. Overview of the project area along the Palos Verdes Peninsula showing the urchin barren extent (pink) mapped in 2010. The locations of restoration areas completed in Years 1 through 7 are in green. Areas restored in Year 8 are in blue. (ESRI 2021)

Pre and Post Restoration Urchin Density Maps – July 1, 2020 through June 30, 2021



Map A2. Density of *S. purpuratus* (individuals per square meter) pre-restoration at White Point, Palos Verdes, California. Average *S. purpuratus* density for this site was 16.66 per m², with some localized areas exceeding 150 per m². The area highlighted in green was restored in Year 7.

(ESRI 2019)



Map A3. Density of *S. purpuratus* (individuals per square meter) post-restoration at White Point, Palos Verdes, California. Average *S. purpuratus* density for this site after restoration was 1.76 per m². The area highlighted in green was restored in Year 7. (ESRI 2021)

Appendix B: CRANE Data Tables 2011 – 2021.

Restoration began at the end of 2014 leading into 2015 at the site Marguerite Central (previously a control site) and was completed in the winter of 2016. Marguerite Central is designated as Restoration for the 2017 surveys.

Table B1. CRANE Survey Metadata.

| | | <u>Survey Dates</u> | | | | | | | | | | |
|--------------------|----------------------|--------------------------------|------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Designation | Site | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 |
| Restoration | Underwater Arch Cove | 2/7/11 | 6/12/12 | 6/13/13 | 7/11/14 | 9/23/15 | 6/22/16 | 7/18/17 | 6/22/18 | 6/12/19 | 6/26/20 | Aug-21 |
| | Honeymoon Cove | 1/28/11 | 3/13/12 | 5/31/13 | 7/2/14 | 8/19/15 | 6/22/16 | 7/18/17 | 6/22/18 | 6/12/19 | 6/24/20 | Aug-21 |
| | Hawthorne | 5/3/11 | 6/12/12 | 6/11/13 | 6/19/14 | 10/7/15 | 9/30/16 | 8/25/17 | 7/11/18 | 6/14/19 | 6/26/20 | Aug-21 |
| | Marguerite Central | 5/3/11 | 6/8/12 | 7/3/13 | 6/20/14 | 9/23/15 | 7/26/16 | 7/18/17 | 7/20/18 | 6/28/19 | 7/9/20 | Aug-21 |
| Reference | Ridges North | 8/12/11 | 7/17/12 | 4/26/13 | 10/29/14 | 9/11/15 | 6/3/16 | 6/30/17 | 7/11/18 | 6/12/19 | 7/23/20 | Aug-21 |
| | Rocky Point North | 6/24/11 | 6/29/12 | 7/2/13 | 7/11/14 | 9/25/15 | 6/10/16 | 6/29/17 | 7/6/18 | 6/19/19 | 7/2/20 | Aug-21 |
| | Point Vicente West | 10/12/11 | 8/10/12 | 4/24/13 | 4/18/14 | 9/23/15 | 6/22/16 | 7/25/17 | 7/18/18 | 6/14/19 | 8/14/20 | Aug-21 |
| | | <u>Bottom Temperature (°C)</u> | | | | | | | | | | |
| Designation | Site | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 |
| Restoration | Underwater Arch Cove | 15.0 | 19.0 | 15.0 | 15.8 | 21.5 | 15.0 | 18.5 | 18.0 | 15.5 | 16.0 | NA |
| | Honeymoon Cove | 15.0 | 11.5 | 18.0 | 16.5 | 18.8 | 16.2 | 20.3 | 18.3 | 15.8 | 16.0 | NA |
| | Hawthorne | 14.4 | 19.0 | 17.0 | 17.0 | 21.0 | 18.0 | 16.8 | 20.6 | 16.0 | 15.0 | NA |
| | Marguerite Central | 15.0 | 17.0 | 17.0 | 20.0 | 22.0 | 14.0 | 20.0 | 20.0 | 18.0 | 19.7 | NA |
| Reference | Ridges North | 18.0 | 16.6 | 13.7 | 19.8 | 21.0 | 15.0 | 17.9 | 22.0 | 16.5 | 12.6 | NA |
| | Rocky Point North | 18.0 | 15.0 | 18.0 | 21.0 | 21.0 | 14.3 | 16.8 | 19.5 | 16.5 | 17.0 | NA |
| | Point Vicente West | 11.0 | 19.0 | 13.2 | 13.5 | 21.0 | 15.2 | 19.7 | 19.5 | 16.5 | 16.2 | NA |
| | | <u>Coordinates</u> | | | | | | | | | | |
| Designation | Site | Latitude | Longitude | | | | | | | | | |
| Restoration | Underwater Arch Cove | 33.75291 | -118.41499 | | | | | | | | | |
| | Honeymoon Cove | 33.76459 | -118.42406 | | | | | | | | | |
| | Hawthorne | 33.75068 | -118.41558 | | | | | | | | | |
| | Marguerite Central | 33.75694 | -118.41772 | | | | | | | | | |
| Reference | Ridges North | 33.78697 | -118.42065 | | | | | | | | | |
| | Rocky Point North | 33.77966 | -118.42739 | | | | | | | | | |
| | Point Vicente West | 33.74073 | -118.41283 | | | | | | | | | |

Table B2. Fish Species Richness (total number of species).

| Designation | Site | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 |
|-------------|----------------------|------|------|------|------|------|------|------|------|------|------|------|
| Restoration | Underwater Arch Cove | 6 | 9 | 6 | 12 | 8 | 8 | 11 | 9 | 9 | 9 | 8 |
| | Honeymoon Cove | 0 | 2 | 4 | 8 | 5 | 12 | 7 | 8 | 8 | 5 | 5 |
| | Marguerite Central | 6 | 10 | 10 | 9 | 11 | 11 | 8 | 9 | 12 | 9 | 11 |
| | Hawthorne | 10 | 6 | 8 | 7 | 10 | 13 | 12 | 12 | 12 | 7 | 9 |
| Reference | Ridges North | 6 | 11 | 7 | 6 | 5 | 10 | 5 | 12 | 8 | 7 | 5 |
| | Rocky Point North | 8 | 8 | 8 | 9 | 6 | 7 | 9 | 11 | 8 | 4 | 6 |
| | Point Vicente West | 8 | 6 | 10 | 11 | 12 | 14 | 9 | 11 | 10 | 12 | 8 |

Yellow indicates when the three original restoration sites (Underwater Arch, Honeymoon Cove, and Hawthorne Restoration) were completed. Marguerite Central, initially a control site, was converted to a restoration site and was completed in 2016.

Table B3. Density of kelp, understory algal species, and invertebrates (individuals per 100 meters squared).

| Species | Designation | Site | #/100 m ² ± SE | | | | | | | | | | |
|----------------------------------|-------------|----------------------|---------------------------|--------------|--------------|-------------|------------|--------------|------------|-----------|------------|------------|-------------|
| | | | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 |
| <i>Anthopleura sola</i> | Restoration | Underwater Arch Cove | 52.5 ± 5.8 | 115 ± 51.7 | 24.2 ± 10.8 | 18.3 ± 5 | 4.2 ± 2.5 | 29.6 ± 7.5 | 11.7 ± 10 | 2.1 ± 0.8 | 3.3 ± 2.4 | — | 0.83 ± 0.83 |
| | | Honeymoon Cove | — | 5 ± 1.7 | — | 1.7 ± 1.7 | — | 2.5 ± 1.6 | — | 0.8 ± 0.8 | — | — | — |
| | | Hawthorne | 418.3 ± 408.3 | 43.3 ± 6.7 | 10 ± 8.3 | 1.7 ± 1.7 | 6.7 ± 5 | 4.2 ± 2.5 | 3.3 ± 3.3 | 8.3 ± 2.4 | 1.7 ± 1 | — | 0.83 ± 0.83 |
| | | Marguerite Central | 79.17 ± 52.5 | 34.17 ± 20.8 | 85.83 ± 9.2 | 33.33 ± 10. | 16.67 ± .0 | 36.67 ± 15.9 | 7.5 ± .8 | — | — | — | — |
| | Reference | Ridges North | — | 1.7 ± 1.7 | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | 0.8 ± 0.8 | — | — | — | — | — | — | — | — | — | 0.83 ± 0.83 |
| | | Point Vicente West | 85.8 ± 4.2 | 155.8 ± 77.5 | 144.2 ± 44.2 | 198.3 ± 6.7 | 0.8 ± 0.8 | 5.8 ± 1.7 | — | 1.7 ± 1.2 | 0.4 ± 0.4 | 6.67 ± 5.0 | — |
| <i>Aplysia californica</i> | Restoration | Underwater Arch Cove | — | — | — | 4.2 ± 2.5 | — | 0.4 ± 0.4 | 10.8 ± 7.5 | — | 0.4 ± 0.4 | — | — |
| | | Honeymoon Cove | — | 0.8 ± 0.8 | 0.8 ± 0.8 | 0.8 ± 0.8 | — | — | — | — | — | — | — |
| | | Hawthorne | 0.8 ± 0.8 | — | 0.8 ± 0.8 | 1.7 ± 1.7 | — | 2.5 ± 2.5 | — | — | — | — | — |
| | | Marguerite Central | 4.17 ± 4.2 | — | — | — | — | 3.75 ± 2.7 | — | — | — | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | 0.8 ± 0.5 | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Aplysia vaccaria</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | 6.3 ± 4 | — | 0.8 ± 0.8 | 1.7 ± 1.7 | — | 2.5 ± 2.5 |
| | | Honeymoon Cove | — | — | — | 0.8 ± 0.8 | — | 3.3 ± 1 | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | 5.8 ± 2.5 | — | — | — | — |
| | | Marguerite Central | — | — | — | — | — | — | — | — | 5.83 ± 5.8 | — | — |
| | Reference | Ridges North | — | — | — | — | — | 0.8 ± 0.8 | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | 1.3 ± 1.3 | — | — | — | — | — |
| <i>Arbacia incisa</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | 0.4 ± 0.4 | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Centrostephanus coronatus</i> | Restoration | Underwater Arch Cove | — | — | — | — | 0.8 ± 0.8 | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Hawthorne | — | 0.8 ± 0.8 | — | — | — | — | 0.8 ± 0.8 | — | — | — | — |

| | | | | | | | | | | | | | | |
|------------------------------------|-------------|----------------------|-------------|--------------|-------------|---------------|--------------|--------------|---------------|--------------|---------------|---------------|---|---|
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — | |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | |
| | | Point Vicente West | — | 0.8 ± 0.8 | — | — | — | — | — | — | — | — | — | |
| <i>Linckia columbiae</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | 0.4 ± 0.4 | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | 2.5 ± 2.5 | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — | — |
| <i>Lytechinus pictus</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | 0.8 ± 0.8 | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — | — |
| <i>Macrocystis pyrifera</i> | Restoration | Underwater Arch Cove | — | — | — | 25 ± 0 | 45.8 ± 4.2 | — | 39.2 ± 5.8 | 95 ± 14.4 | 3.3 ± 2.3 | 37.5 ± .8 | — | |
| | | Honeymoon Cove | 1.7 ± 1.7 | — | — | 7.5 ± 7.5 | 118.3 ± 0 | 88.8 ± 40.4 | 110.8 ± 25.8 | 105.4 ± 11.3 | 150.4 ± 12.5 | 56.67 ± 1.7 | — | |
| | | Hawthorne | 3.3 ± 3.3 | — | — | 3.3 ± 3.3 | 56.7 ± 3.3 | 1.7 ± 1.7 | 158.3 ± 20 | 65.4 ± 7.7 | 30.8 ± 7.1 | 32.5 ± 10.8 | — | |
| | | Marguerite Central | — | — | — | 10.0 ± 10.0 | 9.17 ± 4.2 | 25.0 ± 19. | 95.0 ± 8.3 | 35.83 ± 2.5 | 130.0 ± 40. | 38.33 ± 10.0 | — | |
| | Reference | Ridges North | 67.5 ± 15.8 | 151.7 ± 13.3 | 69.2 ± 22.5 | 87.5 ± 4.2 | 37.5 ± 12.5 | 13.3 ± 3.3 | 78.3 ± 16.7 | 42.5 ± 6 | 117.9 ± 19.4 | 27.5 ± 2.5 | — | |
| | | Rocky Point North | 110 ± 15 | 20 ± 3.3 | 76.7 ± 15 | 319.2 ± 169.2 | 27.5 ± 0.8 | 57.1 ± 35 | 54.2 ± 2.5 | 57.9 ± 7.9 | 87.5 ± 22.1 | 49.17 ± 2.5 | — | |
| | | Point Vicente West | 28.3 ± 6.7 | 27.5 ± 10.8 | 12.5 ± 0.8 | 5.8 ± 2.5 | 13.3 ± 8.3 | 29.6 ± 19.7 | 69.2 ± 12.5 | 41.7 ± 7.6 | 105.8 ± 18.5 | 67.5 ± 7.5 | — | |
| <i>Macrocystis pyrifera Stipes</i> | Restoration | Underwater Arch Cove | — | — | — | 67.5 ± 2.5 | 183.3 ± 10 | — | 162.5 ± 42.5 | 303.8 ± 69.1 | 9.2 ± 6.5 | 166.67 ± 1.7 | — | |
| | | Honeymoon Cove | 3.3 ± 3.3 | — | — | 25.8 ± 25.8 | 564.2 ± 15.8 | 312.5 ± 168 | 481.7 ± 75 | 680.8 ± 65 | 499.2 ± 47 | 666.67 ± 68.3 | — | |
| | | Hawthorne | 31.7 ± 31.7 | — | — | 6.7 ± 6.7 | 354.2 ± 25.8 | 6.7 ± 6.7 | 839.2 ± 232.5 | 612.1 ± 34.3 | 94.2 ± 22.6 | 152.5 ± 42.5 | — | |
| | | Marguerite Central | — | — | — | 28.3 ± 28.3 | 52.5 ± 30.8 | 96.67 ± 76.9 | 470.8 ± 137.5 | 335.8 ± 54.2 | 410.8 ± 182.5 | 607.5 ± 325.8 | — | |

| | | | | | | | | | | | | | |
|----------------------------|-------------|----------------------|----------------|-------------|---------------|----------------|--------------|---------------|--------------|---------------|--------------|---------------|------------|
| | Reference | Ridges North | 830.8 ± 315.8 | 590 ± 186.7 | 531.7 ± 70 | 280 ± 6.7 | 300 ± 55 | 32.5 ± 12.5 | 550 ± 191.7 | 506.3 ± 88.6 | 347.5 ± 61 | 338.33 ± 13.3 | — |
| | | Rocky Point North | 1084.2 ± 107.5 | 235 ± 20 | 986.7 ± 196.7 | 1097.5 ± 619.2 | 157.5 ± 19.2 | 182.1 ± 120.5 | 544.2 ± 19.2 | 621.3 ± 131.9 | 315 ± 109.1 | 646.67 ± 38.3 | — |
| | | Point Vicente West | 449.2 ± 219.2 | 195 ± 36.7 | 136.7 ± 8.3 | 35 ± 18.3 | 159.2 ± 89.2 | 147.1 ± 112.5 | 585.8 ± 69.2 | 363.8 ± 59.9 | 388.3 ± 56.1 | 772.5 ± 109.2 | — |
| <i>Megastraea undosa</i> | Restoration | Underwater Arch Cove | — | — | — | 0.8 ± 0.8 | 5 ± 5 | 31.3 ± 5.6 | 13.3 ± 13.3 | 66.3 ± 17.8 | 53.3 ± 14.7 | 24.17 ± 5.8 | — |
| | | Honeymoon Cove | 2.5 ± 0.8 | — | 0.8 ± 0.8 | 0.8 ± 0.8 | 9.2 ± 0.8 | 5 ± 3.9 | 26.7 ± 8.3 | 112.9 ± 24.5 | 15.4 ± 8.9 | 50.0 ± 1.7 | — |
| | | Hawthorne | 0.8 ± 0.8 | — | — | 0.8 ± 0.8 | 25 ± 25 | 15.8 ± 5.8 | 52.5 ± 9.2 | 99.2 ± 15 | 44.2 ± 8.3 | 11.67 ± 6.7 | — |
| | | Marguerite Central | — | — | — | — | .83 ± .8 | 74.17 ± 32.9 | 50.83 ± 22.5 | 90.83 ± 32.5 | 57.5 ± 12.5 | 165.0 ± 96.7 | — |
| | Reference | Ridges North | 15 ± 13.3 | 6.7 ± 0 | 6.7 ± 5 | — | — | — | — | 1.7 ± 1.2 | 1.3 ± 0.4 | — | 1.7 ± 1.7 |
| | | Rocky Point North | 27.5 ± 9.2 | 25.8 ± 4.2 | 4.2 ± 0.8 | 0.8 ± 0.8 | 3.3 ± 0 | 1.3 ± 1.3 | 0.8 ± 0.8 | 2.1 ± 1.3 | 0.8 ± 0.5 | .83 ± .8 | 0.83 ± 0.8 |
| | | Point Vicente West | — | — | 0.8 ± 0.8 | — | 1.7 ± 0 | 0.8 ± 0.8 | 3.3 ± 1.7 | 5.4 ± 1.4 | 18.3 ± 5.6 | 47.5 ± 19.2 | — |
| <i>Megathura crenulata</i> | Restoration | Underwater Arch Cove | 0.8 ± 0.8 | 2.5 ± 0.8 | 1.7 ± 1.7 | 0.8 ± 0.8 | 0.8 ± 0.8 | — | — | — | — | — | — |
| | | Honeymoon Cove | — | 1.7 ± 0 | — | 0.8 ± 0.8 | — | 0.8 ± 0.8 | — | — | — | — | — |
| | | Hawthorne | 7.5 ± 0.8 | 5 ± 1.7 | 5 ± 1.7 | 3.3 ± 1.7 | — | — | 1.7 ± 1.7 | — | — | — | — |
| | | Marguerite Central | 1.67 ± 1.7 | 1.67 ± 1.7 | 1.67 ± 0.0 | 1.67 ± 1.7 | .83 ± .8 | 2.08 ± 1.6 | — | — | — | — | — |
| | Reference | Ridges North | — | — | 0.8 ± 0.8 | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | 2.5 ± 2.5 | — | 0.8 ± 0.8 | — | — | — | — | — | — | — |
| | | Point Vicente West | 17.5 ± 2.5 | 6.7 ± 1.7 | 6.7 ± 5 | 8.3 ± 1.7 | — | — | — | — | — | — | — |
| <i>Muricea californica</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — |
| | | Marguerite Central | — | .83 ± .8 | — | — | — | — | — | — | — | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | 0.8 ± 0.8 | — | — | — | — | — | — | — |
| <i>Neobernaya spadicea</i> | Restoration | Underwater Arch Cove | 1.7 ± 1.7 | — | — | — | — | — | — | 0.4 ± 0.4 | 0.4 ± 0.4 | — | — |
| | | Honeymoon Cove | — | — | — | 1.7 ± 1.7 | — | 0.8 ± 0.8 | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | — | — | 0.4 ± 0.4 | — | — |
| | | Marguerite Central | — | — | — | — | — | .42 ± .4 | — | — | — | — | — |
| | Reference | Ridges North | — | — | 2.5 ± 2.5 | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | 0.4 ± 0.4 | — | — | 0.4 ± 0.4 | — | — |
| <i>Norrisia norrisi</i> | Restoration | Underwater Arch Cove | 0.8 ± 0.8 | 0.8 ± 0.8 | 0.8 ± 0.8 | — | — | — | — | — | — | — | |

| | | | | | | | | | | | | | | |
|--|-------------|----------------------|----------------|---------------|--------------|-------------|-------------|-------------|--------------|---------------|---------------|--------------|------------|---|
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | 3.75 ± 2.8 | — | — | .83 ± .8 | — | |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | 0.4 ± 0.4 | — | |
| | | Rocky Point North | — | — | — | — | — | — | 5 ± 2.5 | — | — | 0.4 ± 0.4 | .83 ± .8 | |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — | |
| <i>Sargassum sp</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | — | — | — | — | 0.8 ± 0.8 | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| Point Vicente West | | — | — | — | — | — | — | — | — | — | — | — | — | |
| <i>Stephanocystis dioica</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | 13.33 ± .0 | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | 73.8 ± 54.3 | 5.83 ± 4.2 | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | 3.33 ± 3.3 | |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| Point Vicente West | | — | — | — | — | — | — | — | — | — | 6.7 ± 3.6 | — | — | |
| <i>Stephanocystis osmundacea</i> | Restoration | Underwater Arch Cove | — | — | 11.7 ± 1.7 | 10.8 ± 4.2 | 22.5 ± 22.5 | 0.4 ± 0.4 | 3.3 ± 3.3 | 9.2 ± 7 | 27.1 ± 10.1 | — | — | |
| | | Honeymoon Cove | 2.5 ± 2.5 | — | — | — | 0.8 ± 0.8 | — | 2.5 ± 2.5 | 8.8 ± 2.6 | 47.9 ± 6.1 | 20.83 ± 14.2 | — | |
| | | Hawthorne | 6.7 ± 6.7 | 3.3 ± 3.3 | — | 3.3 ± 1.7 | 9.2 ± 9.2 | — | 0.8 ± 0.8 | 9.6 ± 4.7 | 5.8 ± 2.9 | 2.5 ± .8 | — | |
| | | Marguerite Central | — | .83 ± .8 | — | — | — | — | 80.83 ± 73.8 | 6.67 ± 1.7 | 4.17 ± 2.5 | 12.5 ± .8 | 3.33 ± 1.7 | |
| | Reference | Ridges North | — | — | — | — | 10 ± 8.3 | 40 ± 6.7 | 3.3 ± 1.7 | 7.1 ± 4.5 | 12.9 ± 4.4 | — | — | |
| | | Rocky Point North | — | — | — | 2.5 ± 0.8 | 17.5 ± 17.5 | 21.7 ± 12.8 | 4.2 ± 0.8 | 10 ± 2.7 | 20 ± 5 | — | 2.5 ± 2.5 | |
| Point Vicente West | | 0.8 ± 0.8 | — | — | — | 1.7 ± 1.7 | — | 0.8 ± 0.8 | 6.7 ± 6.1 | — | — | — | | |
| <i>Strongylocentrotus franciscanus</i> | Restoration | Underwater Arch Cove | 54.2 ± 17.5 | 33.3 ± 13.3 | 23.3 ± 18.3 | 42.5 ± 7.5 | 2.5 ± 0.8 | 1.3 ± 0.4 | 5.8 ± 0.8 | 10 ± 2.5 | 2.9 ± 1.4 | 2.5 ± 2.5 | — | |
| | | Honeymoon Cove | 63.3 ± 1.7 | 44.2 ± 0.8 | 34.2 ± 4.2 | 11.7 ± 1.7 | 7.5 ± 2.5 | 1.3 ± 0.8 | 0.8 ± 0.8 | 2.1 ± 1 | 2.1 ± 0.8 | 3.33 ± .0 | 0.83 ± 0.8 | |
| | | Hawthorne | 110.8 ± 85.8 | 106.7 ± 6.7 | 51.7 ± 0 | 37.5 ± 10.8 | 2.5 ± 0.8 | 0.8 ± 0.8 | 9.2 ± 4.2 | 6.7 ± 3.6 | 6.7 ± 5.6 | 1.67 ± .0 | — | |
| | | Marguerite Central | 45.0 ± 16.7 | 58.3 ± 30.0 | 12.5 ± 0.8 | 8.3 ± 3.3 | 2.5 ± 2.5 | 1.25 ± .8 | — | 2.5 ± 0.8 | 3.3 ± 3.3 | — | 1.67 ± 0 | |
| | Reference | Ridges North | 2.5 ± 0.8 | 37.5 ± 37.5 | 7.5 ± 0.8 | 4.2 ± 0.8 | 0.8 ± 0.8 | — | — | — | 0.4 ± 0.4 | .83 ± .8 | — | |
| | | Rocky Point North | 5 ± 5 | 9.2 ± 9.2 | 1.7 ± 1.7 | 0.8 ± 0.8 | — | — | — | — | — | — | — | |
| Point Vicente West | | 31.7 ± 10 | 55.8 ± 27.5 | 32.5 ± 4.2 | 26.7 ± 10 | 2.5 ± 0.8 | 2.1 ± 0.8 | — | — | 2.5 ± 1.4 | 1.67 ± .0 | 0.83 ± 0.8 | | |
| <i>Strongylocentrotus purpuratus</i> | Restoration | Underwater Arch Cove | 2195.8 ± 470.8 | 939.2 ± 349.2 | 1008.3 ± 465 | 24.2 ± 10.8 | 1.7 ± 0 | 2.1 ± 1.6 | 13.3 ± 5 | 299.6 ± 141.4 | 221.7 ± 115.4 | 177.5 ± 4.2 | — | |

Table B4. Fish Density (individuals per 100 meters squared).

| Species | Designation | Site | #/100 m ² ± SE | | | | | | | | | | | |
|-------------------------------|-------------|----------------------|---------------------------|-----------|-----------|-----------|------------|-------------|-------------|-------------|-----------|-----------|-------------|-----------|
| | | | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | |
| <i>Anisotremus davidsonii</i> | Restoration | Underwater Arch Cove | — | — | — | — | 0.4 ± 0.4 | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | 0.4 ± 0.3 | 2.5 ± 2.5 | — | — | — | — | |
| | | Hawthorne | — | — | — | — | 0.8 ± 0.8 | 5.8 ± 4.2 | — | 0.2 ± 0.2 | 0.2 ± 0.2 | — | — | |
| | | Marguerite Central | — | — | — | — | 27.9±20.9 | .6±.6 | 1.3±1.3 | — | — | .8±.8 | — | |
| | Reference | Ridges North | — | — | — | — | — | 0.4 ± 0.4 | — | 9.4 ± 6 | 0.2 ± 0.2 | — | 9.17 ± 8.1 | |
| | | Rocky Point North | — | — | — | — | — | — | 5.4 ± 4.9 | 8.8 ± 6 | 0.2 ± 0.2 | — | — | |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — | |
| Atherinopsidae | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | 54.6 ± 47.6 | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | 5.4±5.4 | — | .4±.4 | — | 22.9±15.7 | — | — | |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — | |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | |
| | | Point Vicente West | — | — | — | — | — | 0.2 ± 0.2 | — | — | — | — | — | |
| <i>Brachyistius frenatus</i> | Restoration | Underwater Arch Cove | — | — | — | 1.3 ± 1.3 | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | 0.8 ± 0.8 | 0.2 ± 0.2 | — | — | 0.8 ± 0.5 | |
| | | Hawthorne | — | — | — | — | — | — | — | 2.7 ± 1.5 | — | — | 1.25 ± 0.8 | |
| | | Marguerite Central | — | — | — | — | — | — | — | 12.1±10.0 | — | 1.7±.7 | 7.5 ± 6.9 | |
| | Reference | Ridges North | 0.4 ± 0.4 | 0.4 ± 0.4 | — | — | — | — | — | — | — | 2.5±1.1 | — | |
| | | Rocky Point North | 1.7 ± 0.7 | 0.4 ± 0.4 | 0.8 ± 0.8 | 0.4 ± 0.4 | — | — | 0.4 ± 0.4 | 2.5 ± 0.6 | 0.2 ± 0.2 | .4±.4 | 0.8 ± 0.5 | |
| | | Point Vicente West | 0.4 ± 0.4 | 2.5 ± 1.6 | 0.8 ± 0.5 | — | 14.2 ± 5.9 | 0.8 ± 0.8 | 33.8 ± 16.9 | 29.6 ± 17.9 | — | .4±.4 | 44.6 ± 33.2 | |
| <i>Chromis punctipinnis</i> | Restoration | Underwater Arch Cove | 1.3 ± 1.3 | 1.3 ± 1.3 | 0.4 ± 0.4 | 2.1 ± 0.8 | — | 10.6 ± 5.6 | 2.5 ± 2.5 | 0.6 ± 0.3 | 0.4 ± 0.3 | — | 0.42 ± 0.4 | |
| | | Honeymoon Cove | — | — | — | — | — | 1.3 ± 1.3 | — | — | — | — | — | |
| | | Hawthorne | 0.4 ± 0.4 | — | — | — | — | — | 10.4 ± 10.4 | — | — | 0.4 ± 0.3 | — | 2.9 ± 2.9 |
| | | Marguerite Central | .8±.8 | — | 6.3±5.2 | — | — | — | — | — | 20.4±18.3 | — | 41.7 ± 41.7 | |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | 1.7±1.7 | — | |
| | | Rocky Point North | 0.4 ± 0.4 | — | — | — | — | — | — | 0.4 ± 0.4 | — | — | — | |
| | | Point Vicente West | — | — | — | 8.3 ± 5.5 | — | 12.7 ± 10.6 | — | 0.2 ± 0.2 | 1.7 ± 0.8 | 41.3±15.5 | 1.25 ± 0.8 | |
| <i>Damalichthys vacca</i> | Restoration | Underwater Arch Cove | — | — | — | 0.4 ± 0.4 | — | — | — | 0.2 ± 0.2 | — | — | 0.42 ± 0.4 | |
| | | Honeymoon Cove | — | — | — | 0.4 ± 0.4 | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | 0.4 ± 0.4 | — | — | — | — | |
| | | Marguerite Central | .8±.5 | .8±.5 | — | 3.3±3.3 | — | — | — | — | — | — | 1.25 ± 1.25 | |
| | Reference | Ridges North | — | — | — | 0.4 ± 0.4 | — | — | — | — | — | .4±.4 | — | |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | |
| | | Point Vicente West | — | 0.8 ± 0.5 | — | 0.4 ± 0.4 | — | — | — | — | — | — | — | |
| <i>Embiotoca jacksoni</i> | Restoration | Underwater Arch Cove | 2.9 ± 2.4 | 0.4 ± 0.4 | 0.4 ± 0.4 | 0.8 ± 0.8 | 0.8 ± 0.5 | 0.2 ± 0.2 | 0.4 ± 0.4 | 3.1 ± 0.7 | 0.6 ± 0.4 | 1.3±.4 | 0.8 ± 0.5 | |
| | | Honeymoon Cove | — | 0.4 ± 0.4 | — | — | — | 0.4 ± 0.3 | — | 1 ± 0.5 | 0.2 ± 0.2 | .4±.4 | 0.4 ± 0.4 | |
| | | Hawthorne | 3.8 ± 0.4 | 2.5 ± 1.1 | 2.1 ± 0.4 | 2.9 ± 1.3 | 8.3 ± 8.3 | 1.3 ± 1.3 | 2.5 ± 1.1 | — | 2.9 ± 1.7 | 2.1±1.6 | 0.4 ± 0.4 | |
| | | Marguerite Central | .4±.4 | 2.1±.8 | 1.7±.7 | 2.1±1.3 | 1.3±.8 | .8±.3 | 2.1±1.3 | — | 3.3±1.5 | 1.7±1.0 | 1.25 ± 0.4 | |

| | | | | | | | | | | | | | | |
|--------------------------------|-------------|----------------------|-----------|-----------|-----------|-----------|-----------|------------|-------------|------------|-----------|---------|------------|-----------|
| | Reference | Ridges North | 0.4 ± 0.4 | — | — | — | — | 0.4 ± 0.4 | — | 0.4 ± 0.3 | 0.8 ± 0.5 | — | — | |
| | | Rocky Point North | 2.9 ± 1.4 | 3.8 ± 1.7 | 0.4 ± 0.4 | 0.8 ± 0.5 | 0.8 ± 0.5 | 1.7 ± 1.4 | — | 0.8 ± 0.4 | 0.6 ± 0.3 | — | — | |
| | | Point Vicente West | 1.7 ± 0.7 | 0.4 ± 0.4 | 6.3 ± 0.8 | 2.5 ± 1.1 | 1.3 ± 0.8 | 1.7 ± 0.8 | — | 0.2 ± 0.2 | 1.9 ± 0.6 | 7.5±6.4 | 0.8 ± 0.5 | |
| <i>Gibbonsia elegans</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | 0.4 ± 0.4 | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | 0.4 ± 0.4 | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — | — |
| <i>Girella nigricans</i> | Restoration | Underwater Arch Cove | 4.2 ± 3.6 | 0.4 ± 0.4 | — | 0.4 ± 0.4 | — | 0.2 ± 0.2 | 0.8 ± 0.5 | 0.8 ± 0.3 | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | 0.8 ± 0.8 | 0.6 ± 0.4 | — | — | — | — | — | |
| | | Hawthorne | 2.1 ± 0.8 | 2.5 ± 1.6 | 0.8 ± 0.5 | — | 1.7 ± 1.7 | 0.4 ± 0.4 | 1.7 ± 1.2 | — | 1.9 ± 1.6 | — | — | |
| | | Marguerite Central | — | 1.7±1.7 | — | .4±.4 | 20.±13.1 | 18.5±8.1 | .4±.4 | .8±.8 | 3.8±1.7 | 1.3±1.3 | 15 ± 9.3 | |
| | Reference | Ridges North | — | — | — | — | — | — | — | 0.4 ± 0.3 | 0.2 ± 0.2 | — | — | — |
| | | Rocky Point North | — | 0.8 ± 0.8 | — | 1.7 ± 1 | 1.3 ± 0.8 | 2.1 ± 1.2 | — | — | 0.2 ± 0.2 | — | — | — |
| | | Point Vicente West | — | — | 1.3 ± 0.8 | 1.7 ± 1.2 | 0.4 ± 0.4 | 0.8 ± 0.3 | 1.3 ± 0.8 | 0.8 ± 0.6 | 1.3 ± 0.5 | 2.1±1.3 | — | — |
| Gobiidae | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | 2.9 ± 2.9 | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | — | 4.2 ± 4.2 | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | 12.5 ± 8.8 | — | — | — | — | — | — |
| <i>Haemulon californiensis</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — | |
| | Reference | Ridges North | — | — | — | — | — | — | — | 7.7 ± 7.2 | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | 8.3 ± 8.3 | 224 ± 84.5 | — | — | — | 0.8 ± 0.5 |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — | — |
| <i>Halichoeres semicinctus</i> | Restoration | Underwater Arch Cove | 0.8 ± 0.5 | — | 0.4 ± 0.4 | 0.8 ± 0.8 | 2.1 ± 0.8 | 1.9 ± 0.7 | 16.3 ± 11.8 | 3.1 ± 1 | 1.7 ± 0.7 | 1.7±.7 | — | |
| | | Honeymoon Cove | — | — | — | 2.9 ± 1.7 | — | 4.8 ± 0.7 | 6.7 ± 1.2 | 2.3 ± 0.9 | 4.4 ± 1.5 | .8±.8 | 0.83 ± 0.5 | |
| | | Hawthorne | 0.8 ± 0.8 | — | — | — | 3.3 ± 1.5 | 8.8 ± 4.2 | 16.7 ± 11 | 3.5 ± 0.8 | 4.4 ± 1.1 | 5.8±3.8 | 1.25 ± 0.8 | |
| | | Marguerite Central | — | — | — | — | 7.5±1.6 | 18.1±7.2 | — | 7.1±2.4 | 2.1±1.0 | 1.3±.4 | 1.6 ± 0.9 | |
| | Reference | Ridges North | — | 1.3 ± 0.8 | 1.3 ± 0.8 | — | 4.2 ± 0.5 | 2.1 ± 0.8 | 0.4 ± 0.4 | 0.6 ± 0.3 | 0.6 ± 0.3 | 1.3±.8 | — | |
| | | Rocky Point North | 2.1 ± 1.6 | — | 5 ± 2.4 | 0.4 ± 0.4 | 8.3 ± 2 | 1.9 ± 0.9 | 4.2 ± 1.6 | 2.5 ± 1.1 | 1 ± 0.6 | — | 0.42 ± 0.4 | |
| | | Point Vicente West | 0.4 ± 0.4 | — | 0.4 ± 0.4 | — | 1.3 ± 0.4 | 3.3 ± 0.3 | 1.3 ± 0.4 | 1.3 ± 0.5 | 1.3 ± 0.5 | 3.8±1.3 | — | |
| <i>Hermosilla azurea</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | 0.4 ± 0.4 | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | 1.3±.8 | 11.5±7.5 | — | 1.3±.8 | — | — | — | |

| | | | | | | | | | | | | | |
|-------------------------------------|-------------|----------------------|------------|-----------|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Medialuna californiensis</i> | Restoration | Underwater Arch Cove | — | — | — | — | 2.5 ± 1.4 | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | 0.2 ± 0.2 | — | — | — | — | — |
| | | Hawthorne | 0.4 ± 0.4 | — | — | — | 4.6 ± 2.3 | 1.3 ± 0.4 | — | 0.4 ± 0.3 | 0.2 ± 0.2 | — | — |
| | | Marguerite Central | — | .4±.4 | — | — | 4.2±2.5 | .2±.2 | — | — | .8±.5 | .4±.4 | 0.4 ± 0.4 |
| | Reference | Ridges North | — | — | — | — | — | 2.1 ± 1.3 | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | 0.6 ± 0.6 | — | — | — | — | — |
| | | Point Vicente West | 0.4 ± 0.4 | — | — | — | 2.5 ± 0.8 | 1 ± 0.5 | 0.8 ± 0.8 | — | — | .4±.4 | — |
| <i>Oxyjulis californica</i> | Restoration | Underwater Arch Cove | — | 0.4 ± 0.4 | 0.8 ± 0.8 | 1.7 ± 1.2 | — | — | — | — | 1 ± 0.8 | 1.7±1.7 | 2.1 ± 2.1 |
| | | Honeymoon Cove | — | — | — | 0.8 ± 0.8 | 2.9 ± 1.3 | 0.2 ± 0.2 | 12.1 ± 5.6 | — | — | — | — |
| | | Hawthorne | 5 ± 3.4 | 1.3 ± 0.8 | 1.7 ± 1.2 | 2.1 ± 1.6 | 1.7 ± 0.7 | 2.9 ± 1 | — | 0.2 ± 0.2 | 0.6 ± 0.4 | 31.3±30.7 | 2.1 ± 1.2 |
| | | Marguerite Central | — | .8±.8 | 2.9±1.8 | 3.8±2.2 | 5.4±4.9 | 1.5±.9 | — | .4±.4 | 7.1±3.3 | — | 0.8 ± 0.8 |
| | Reference | Ridges North | — | — | — | 0.8 ± 0.8 | — | — | — | 0.2 ± 0.2 | — | — | — |
| | | Rocky Point North | 0.4 ± 0.4 | 0.8 ± 0.5 | — | 0.4 ± 0.4 | — | — | — | — | — | — | — |
| | | Point Vicente West | 13.3 ± 6.7 | 2.1 ± 1 | 1.3 ± 0.4 | 23.3 ± 7.9 | — | — | 8.8 ± 8.2 | — | 0.6 ± 0.4 | 2.5±1.4 | 1.67 ± 1.2 |
| <i>Oxylebius pictus</i> | Restoration | Underwater Arch Cove | — | 0.4 ± 0.4 | — | — | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | 0.4 ± 0.4 | — | 0.4 ± 0.4 | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | 0.4 ± 0.4 | 0.8 ± 0.8 | — | — | — | — | — | — | — |
| | | Marguerite Central | .4±.4 | — | 2.9±1.7 | .4±.4 | — | — | — | — | .4±.4 | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | 0.4 ± 0.4 | 0.4 ± 0.4 | — | 1.3 ± 0.4 | — | — | — | — | — | — | — |
| <i>Paralabrax clathratus</i> | Restoration | Underwater Arch Cove | — | 0.4 ± 0.4 | 0.4 ± 0.4 | 6.7 ± 4.5 | 4.2 ± 2.2 | 3.8 ± 0.8 | 17.1 ± 8.3 | 5.6 ± 1.4 | 2.7 ± 0.7 | 3.8±1 | 5.4 ± 2.1 |
| | | Honeymoon Cove | — | — | 0.8 ± 0.5 | 1.3 ± 0.4 | 0.8 ± 0.8 | 1.3 ± 0.4 | 4.2 ± 3.6 | 1.9 ± 0.6 | 1.5 ± 0.7 | 2.9±1.4 | 0.83 ± 0.8 |
| | | Hawthorne | — | — | 0.8 ± 0.5 | 0.8 ± 0.5 | 0.8 ± 0.5 | 5.4 ± 1.3 | 6.7 ± 3.2 | 3.8 ± 0.4 | 1.3 ± 0.5 | .4±.4 | 4.2 ± 0.8 |
| | | Marguerite Central | — | 1.3 ± 1.3 | 1.7 ± 1.2 | 3.3 ± 0.7 | 5.0 ± 3.0 | 7.1 ± 3.8 | 11.3 ± 0.8 | 9.2 ± 1.0 | 6.7 ± 0.7 | 2.9±1.3 | 3.8 ± 0.8 |
| | Reference | Ridges North | 2.9 ± 0.8 | 0.4 ± 0.4 | 2.1 ± 0.4 | 2.9 ± 0.8 | 2.9 ± 1.4 | 1.7 ± 1.2 | 1.7 ± 1.2 | 4.6 ± 1.2 | — | .4±.4 | 0.8 ± 0.5 |
| | | Rocky Point North | 1.7 ± 0.7 | 2.5 ± 1.4 | 6.7 ± 1.9 | 2.1 ± 0.8 | 4.2 ± 1.6 | 1.5 ± 0.8 | 2.5 ± 1.6 | 2.7 ± 0.5 | 0.4 ± 0.3 | 1.7±.7 | 0.8 ± 0.5 |
| | | Point Vicente West | — | — | 0.8 ± 0.5 | — | 2.9 ± 0.4 | 2.9 ± 1.1 | 7.5 ± 1.1 | 5.2 ± 1.2 | 1.9 ± 0.5 | 7.5±2.0 | 1.7 ± 0.5 |
| <i>Paralabrax maculatofasciatus</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | 0.4 ± 0.4 | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Paralabrax nebulifer</i> | Restoration | Underwater Arch Cove | — | — | — | 1.3 ± 0.8 | — | 0.4 ± 0.3 | 1.3 ± 0.8 | 0.2 ± 0.2 | 0.4 ± 0.4 | .4±.4 | 2.5 ± 1.6 |
| | | Honeymoon Cove | — | — | — | 0.8 ± 0.5 | — | — | — | 1.5 ± 0.6 | 0.2 ± 0.2 | .4±.4 | 0.8 ± 0.8 |
| | | Hawthorne | — | 0.4 ± 0.4 | — | 0.8 ± 0.5 | — | 0.4 ± 0.4 | 2.1 ± 0.8 | 0.6 ± 0.4 | 0.4 ± 0.3 | .4±.4 | — |
| | | Marguerite Central | .4±.4 | — | .4±.4 | 1.3±.8 | — | 1.5±.7 | .8±.5 | — | .8±.8 | — | — |

| | | | | | | | | | | | | | |
|-----------|--------------------|---|---|---|---|---|---|---|-----------|---|---|---|---|
| Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — | — |
| | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| | Point Vicente West | — | — | — | — | — | — | — | 0.2 ± 0.2 | — | — | — | — |

Table B5. Fish Biomass (individuals per 100 meters squared).

| Species | Designation | Site | <i>g/100 m² ± SE</i> | | | | | | | | | | |
|-------------------------------|-------------|----------------------|---------------------------------|-------------|-------------|-------------|---------------|---------------|-----------------|---------------|-------------|-------------|-------------|
| | | | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 |
| <i>Anisotremus davidsonii</i> | Restoration | Underwater Arch Cove | — | — | — | — | 50.5 ± 50.5 | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | 19.9 ± 13 | 119.4 ± 119.4 | — | — | — | — |
| | | Hawthorne | — | — | — | — | 101 ± 101 | 174.7 ± 126.5 | — | 10 ± 10 | 7.9 ± 7.9 | — | — |
| | | Marguerite Central | — | — | — | — | 11117.8±8614. | — | 28.8±28.8 | — | — | 235.5±235.5 | — |
| | Reference | Ridges North | — | — | — | — | — | 5.3 ± 5.3 | — | 532.2 ± 331.3 | 10 ± 10 | — | 1272 ± 1115 |
| | | Rocky Point North | — | — | — | — | — | — | 1154.1 ± 1127.7 | 973.9 ± 521 | 4.8 ± 4.8 | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| Atherinopsidae | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | 995.2 ± 930.8 | — | — | — | — |
| | | Marguerite Central | — | — | — | — | 130.2±130.2 | 78.5±78.5 | 94.5±94.5 | — | 550.9±378.1 | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | 4.1 ± 4.1 | — | — | — | — | — | — | — |
| Point Vicente West | | — | — | — | — | — | 12.7 ± 12.7 | — | — | — | .5±.5 | — | |
| <i>Brachyistius frenatus</i> | Restoration | Underwater Arch Cove | — | — | — | 31.9 ± 31.9 | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | 3.6 ± 2.2 | 5.3 ± 5.3 | — | — | 19.4 ± 16.8 |
| | | Hawthorne | — | — | — | — | — | — | — | 15.1 ± 9.7 | — | — | 18.6 ± 12 |
| | | Marguerite Central | — | — | — | — | — | — | — | 57.4±47.3 | — | 8.2±4.2 | 64.5 ± 51.3 |
| | Reference | Ridges North | 8.1 ± 8.1 | 6 ± 6 | — | — | — | — | — | — | — | 778.4±734.2 | — |
| | | Rocky Point North | 40.9 ± 18.5 | 13.7 ± 13.7 | 12.4 ± 12.4 | 10.6 ± 10.6 | — | — | 1.2 ± 1.2 | 19.6 ± 4.2 | 1 ± 1 | 6.±6.0 | 14 ± 8.3 |
| | | Point Vicente West | 10.6 ± 10.6 | 26.8 ± 17.1 | 11 ± 7.6 | — | 135.3 ± 62.5 | 29.3 ± 29.3 | 217.6 ± 123.9 | 177.8 ± 107.7 | — | 8.1±8.1 | 351.3 ± 235 |
| <i>Chromis punctipinnis</i> | Restoration | Underwater Arch Cove | 60.3 ± 60.3 | 39.1 ± 39.1 | 13 ± 13 | 79.7 ± 39.7 | 8.6 ± 8.6 | 245.6 ± 134.6 | 120.6 ± 120.6 | 25.2 ± 13 | 16.6 ± 11.2 | — | 10.2 ± 10.2 |
| | | Honeymoon Cove | — | — | — | — | — | 49.4 ± 49.4 | — | — | — | — | — |
| | | Hawthorne | 20.1 ± 20.1 | — | 8 ± 5.7 | — | 27.3 ± 27.3 | 282.9 ± 247.7 | 5.1 ± 5.1 | — | 13.3 ± 9 | — | 54.4 ± 54.4 |
| | | Marguerite Central | 20.5±20.5 | 3.0±3.0 | 194.9±153.7 | — | — | — | — | — | 634.4±531.5 | — | 1305 ± 1305 |
| | Reference | Ridges North | — | — | — | — | — | 57.3 ± 33.2 | — | — | — | 85.4±69.6 | — |
| | | Rocky Point North | 10.2 ± 10.2 | — | — | — | — | — | — | 10.2 ± 10.2 | — | — | — |

| | | | | | | | | | | | | | | |
|---------------------------|-------------|----------------------|------------------|------------------|------------------|------------------|--------------------|------------------|-------------------|-----------------|------------------|------------------|------------------|---|
| | | Point Vicente West | — | — | 75.4 ± 75.4 | 290.4 ± 195.3 | 9.4 ± 7.4 | 370.9 ± 319.4 | — | 3 ± 3 | 44.7 ± 23.2 | 1049.5±49 8.5 | 54.8 ± 42.2 | |
| <i>Damalichthys vacca</i> | Restoration | Underwater Arch Cove | — | — | — | 56.6 ± 56.6 | — | — | — | 28.3 ± 28.3 | — | — | 24.3 ± 24.3 | |
| | | Honeymoon Cove | — | — | — | 56.6 ± 56.6 | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | 56.6 ± 56.6 | — | — | — | |
| | | Marguerite Central | 113.2±65.4 | 198.9±182. 4 | — | 819.8±819. 8 | — | — | — | — | — | — | 558.7 ± 558.7 | |
| | Reference | Ridges North | — | — | — | 56.6 ± 56.6 | — | — | — | — | — | — | 12.6±12.6 | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | — |
| Point Vicente West | | — | 116.4 ± 106.8 | — | 109 ± 109 | — | — | — | — | — | — | — | — | |
| <i>Embiotoca jacksoni</i> | Restoration | Underwater Arch Cove | 453.2 ± 371.9 | 14.7 ± 14.7 | 24.2 ± 24.2 | 83.8 ± 83.8 | 42.5 ± 21.6 | 4.3 ± 4.3 | 24.2 ± 24.2 | 213.2 ± 48.4 | 24.3 ± 17.4 | 105.1±45.3 | 44.9 ± 28.8 | |
| | | Honeymoon Cove | — | 64.7 ± 64.7 | — | 2.7 ± 2.7 | — | 18.2 ± 13 | 6.3 ± 5.1 | 66.2 ± 41.6 | 27.2 ± 27.2 | 45.2±45.2 | 70.5 ± 65.2 | |
| | | Hawthorne | 173.3 ± 46.6 | 91.4 ± 34.1 | 266.2 ± 116.2 | 214.9 ± 136.7 | 1294.9 ± 1294.9 | 89.8 ± 89.8 | 126.8 ± 60.3 | 0.8 ± 0.8 | 292.4 ± 219.1 | 206.±177.8 | 45.2 ± 45.2 | |
| | | Marguerite Central | 24.2±24.2 | 192.6±113. 1 | 251.4±114. 7 | 151.2±102. 3 | 138.1±130. 3 | 59.5±32.2 | 97.1±52.8 | 2.7±2.7 | 234.±93.8 | 210.3±128. 2 | 100.2 ± 54.9 | |
| | Reference | Ridges North | 26.9 ± 23.4 | 1.7 ± 1.7 | — | — | — | 64.7 ± 64.7 | — | 13 ± 8.6 | 111.8 ± 80 | — | — | |
| | | Rocky Point North | 313.2 ± 179 | 366.8 ± 153.2 | 138.3 ± 138.3 | 83.8 ± 61.1 | 43.2 ± 25.3 | 55.1 ± 51.8 | — | 61.1 ± 43.8 | 38.5 ± 23.4 | — | — | |
| Point Vicente West | | 78.4 ± 43.5 | 64.7 ± 64.7 | 677.7 ± 183.7 | 474.5 ± 160.9 | 78.9 ± 61.5 | 107.5 ± 58.3 | — | 71.3 ± 70.4 | 159.5 ± 60.6 | 541.9±397. 5 | 32.4 ± 22.8 | | |
| <i>Gibbonsia elegans</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | 6.8 ± 6.8 | — | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — | |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — | |
| | | Marguerite Central | — | — | — | — | — | — | 4693.2±21 99.8 | — | — | — | — | |
| | Reference | Ridges North | — | 8.8 ± 8.8 | — | — | — | — | — | — | — | — | — | |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — | |
| Point Vicente West | | — | — | — | — | — | — | — | — | — | — | — | | |
| <i>Girella nigricans</i> | Restoration | Underwater Arch Cove | 284.7 ± 251.1 | 16.9 ± 16.9 | — | 26.1 ± 26.1 | — | 61.9 ± 61.9 | 52.2 ± 30.1 | 42.6 ± 16.5 | — | — | — | |
| | | Honeymoon Cove | — | — | — | — | 186.6 ± 186.6 | 27.7 ± 21.2 | — | — | — | — | — | |
| | | Hawthorne | 338.1 ± 117.7 | 100.7 ± 59.3 | 186.6 ± 118.7 | — | 863.9 ± 863.9 | 216 ± 216 | 901.4 ± 646.3 | — | 273.7 ± 248.7 | — | — | |
| | | Marguerite Central | — | 238.8±238. 8 | — | 62.7±62.7 | 1161.6±96 6.6 | — | 62.7±62.7 | 125.5±125. 5 | 658.±329.7 | 374.8±374. 8 | 5038.6 ± 3215 | |
| | Reference | Ridges North | — | 3.8 ± 3.8 | — | — | — | — | — | 21.5 ± 14.5 | 31.4 ± 31.4 | — | — | |
| | | Rocky Point North | — | 125.5 ± 125.5 | — | 434.3 ± 255.7 | 463.7 ± 269 | 340.4 ± 185.2 | — | — | 26.8 ± 26.8 | — | — | |


| | | | | | | | | | | | | | |
|---------------------------------|-------------|----------------------|---------------|-------------|--------------|---------------|---------------|--------------|---------------|-------------|-------------|-------------|---------------|
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Lythrypnus dalli</i> | Restoration | Underwater Arch Cove | — | 6.7 ± 6.7 | — | — | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Medialuna californiensis</i> | Restoration | Underwater Arch Cove | — | — | — | — | 236.2 ± 186.7 | — | — | — | — | — | — |
| | | Honeymoon Cove | — | — | — | — | — | 12.4 ± 12.4 | — | — | — | — | — |
| | | Hawthorne | 58.3 ± 58.3 | — | — | — | 846 ± 403.7 | 74.6 ± 24.9 | — | 20.6 ± 13.9 | 29.2 ± 29.2 | — | — |
| | | Marguerite Central | — | 24.9±24.9 | — | — | 404.5±246.8 | 97.0±97.0 | — | — | 32.6±18.8 | 16.3±16.3 | 30.1 ± 30.1 |
| | Reference | Ridges North | — | — | — | — | — | 102.4 ± 59.2 | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | 30.7 ± 30.7 | — | — | — | — | — |
| | | Point Vicente West | 113 ± 113 | — | — | — | 98.2 ± 18.6 | 91.3 ± 47.4 | 171.3 ± 171.3 | — | — | 16.3±16.3 | — |
| <i>Oxyjulis californica</i> | Restoration | Underwater Arch Cove | — | 14.1 ± 14.1 | 10.9 ± 10.9 | 237.3 ± 195.2 | — | — | — | — | 65 ± 62.8 | 313.1±313.1 | 128.3 ± 124.6 |
| | | Honeymoon Cove | — | — | 28.1 ± 28.1 | 28.2 ± 28.2 | 57 ± 19.6 | 7.1 ± 5.6 | 536.3 ± 328.9 | — | 9.5 ± 8 | — | — |
| | | Hawthorne | 142.7 ± 95.1 | 35.2 ± 21.1 | 40.5 ± 31.9 | 63.4 ± 46.5 | 140.1 ± 57.2 | 137.1 ± 62.1 | 11.6 ± 11.6 | 5.7 ± 5.7 | 24.1 ± 15.8 | 495.4±462.5 | 86.2 ± 61.1 |
| | | Marguerite Central | — | 28.2±28.2 | 63.1±36.8 | 147.9±75.8 | 287.9±269.5 | 57.1±33.2 | — | 15.4±15.4 | 347.±137.1 | — | 23.1 ± 23.1 |
| | Reference | Ridges North | — | 0.4 ± 0.4 | 1.3 ± 1.3 | 10.6 ± 10.6 | 1.4 ± 1.2 | — | — | 7.1 ± 7.1 | — | — | — |
| | | Rocky Point North | 3.9 ± 3.9 | 21.1 ± 13.5 | — | 11.4 ± 5.3 | — | — | 1.3 ± 1.3 | — | — | — | — |
| | | Point Vicente West | 862.5 ± 503.5 | 81.3 ± 42.9 | 191.6 ± 70.1 | 700.3 ± 299.7 | 9.2 ± 2.8 | — | 98.4 ± 71.9 | — | 40 ± 28.5 | 109.1±65.2 | 76.2 ± 62 |
| <i>Oxylebius pictus</i> | Restoration | Underwater Arch Cove | — | 14.9 ± 14.9 | — | — | — | — | — | — | — | — | — |
| | | Honeymoon Cove | — | 18.7 ± 18.7 | — | 18.7 ± 18.7 | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | 18.7 ± 18.7 | 38.1 ± 38.1 | — | — | 2.2 ± 2.2 | — | — | — | — |
| | | Marguerite Central | 23.2±23.2 | — | 162.4±95.6 | 23.2±23.2 | — | — | — | — | 14.9±14.9 | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | 23.2 ± 23.2 | 23.2 ± 23.2 | — | 49 ± 17.7 | — | — | — | — | — | — | — |

| | | | | | | | | | | | | | |
|-------------------------|-------------|----------------------|---|-------------|---|---|---|---|---|-------------|-------------|---|---|
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — |
| Reference | | Ridges North | — | 10.7 ± 10.7 | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | — | — | — | — |
| <i>Urobatis halleri</i> | Restoration | Underwater Arch Cove | — | — | — | — | — | — | — | — | 35.7 ± 35.7 | — | — |
| | | Honeymoon Cove | — | — | — | — | — | — | — | — | — | — | — |
| | | Hawthorne | — | — | — | — | — | — | — | — | — | — | — |
| | | Marguerite Central | — | — | — | — | — | — | — | — | — | — | — |
| | Reference | Ridges North | — | — | — | — | — | — | — | — | — | — | — |
| | | Rocky Point North | — | — | — | — | — | — | — | — | — | — | — |
| | | Point Vicente West | — | — | — | — | — | — | — | 28.6 ± 28.6 | — | — | — |

Table B6. White Point CRANE survey data for Year 7.

| White Point Crane Data - September 2020-2021 | | |
|--|---------------|---------------|
| Analysis | 2020 | 2021 |
| Coordinates: | | |
| Latitude | 33.71287 | 33.71287 |
| Longitude | -118.3159 | -118.3159 |
| Temperature (°C) | 17.5 | 16.1 |
| Fish Richness | 10 | 10 |
| Fish Diversity H | 2.029 | 0.48 |
| Fish Diversity 1-D | 0.83 | 0.17 |
| Fish Density: | | |
| <i>Paralabrax clathratus</i> (/100m ²) | 2.1 ± 0.4 | 3.3 ± 2.3 |
| <i>Semicossyphus pulcher</i> (/100m ²) | 1.3 ± 0.4 | 1.3 ± 0.4 |
| Fish Biomass: | | |
| <i>Paralabrax clathratus</i> (g/100m ²) | 373.3 ± 108.5 | 514.3 ± 350 |
| <i>Semicossyphus pulcher</i> (g/100m ²) | 402.2 ± 220.6 | 572.4 ± 572.4 |
| Swath Diversity H | 0.875 | 1.997 |
| Swath Diversity 1-D | 0.35 | 0.832 |
| Swath Density: | | |
| <i>Macrocystis pyrifera stipes</i> (/100m ²) | 11.7 ± 8 | 687.5 ± 60.8 |
| <i>Panulirus interruptus</i> | - | 1.67 ± 1.67 |
| <i>Mesocentrotus franciscanus</i> (/100m ²) | 4.2 ± 4.2 | 15 ± 3.3 |
| <i>Strongylocentrotus purpuratus</i> (/100m ²) | 510 ± 278 | 160.9 ± 6.3 |

Honeymoon Cove



Honeymoon Cove Block T2 (HMC T2) east-west running ridge is a large distinguishable feature easily found by divers. This block was restored in March 2014. GPS: 33.764260, -118.423734



HMC T2 07/29/16




HMC T2 08/07/17



HMC T2 07/18/18



HMC T2 07/18/19



HMC T2 07/17/2020



HMC T2 08/31/21

Honeymoon Cove Block R5 (HMC R5) is the site of another TBF project with ongoing monitoring. Divers visit this area annually to conduct subtidal surveys allowing the opportunity to collect photos over time. This block was restored in November 2014. GPS: 33.765297, -118.424221



HMC R5 06/22/15



HMC R5 09/24/15



HMC R5 11/12/15



HMC R5 02/10/16



HMC R5 08/3/17



HMC R5 07/3/18



HMC R5 07/18/19




HMC R5 07/17/20



HMC R5 08/31/21

Marguerite



Marguerite Block T16 (MARG T16) was monitored monthly by TBF divers for 2 years starting in 2016 for a wave attenuation study. This block was restored in September 2016. Subsequent photo/videos occur annually.
GPS: 33.757561, -118.41782



MARG T16 08/10/16



MARG T16 08/3/17



MARG T16 07/20/18



MARG T16 06/21/19




MARG T16 08/12/20



MARG T16 07/09/21

Underwater Arch Cove




Underwater Arch Cove Blocks (UWAC) J1, J2 and T7 were the locations of our first transect video shot in 2014. In 2016 and 2017, this video transect was recorded again and photos from both dates have been archived. Divers will continue to revisit this area annually for video and photography.

UWAC J1 restoration was complete in November 2013. GPS: 33.75205979, -118.4156861



UWAC J1 08/14/14



UWAC J1 07/07/16



UWAC J1 07/27/17



UWAC J1 07/18/18



UWAC J1 06/21/19



UWAC J1 07/24/20



UWAC J1 07/13/21

UWAC J2 was restored in July 2014. GPS: 33.7523302, -118.4151245



UWAC J2 PRE-RESTORATION 07/12/14



UWAC J2 08/14/14



UWAC J2 07/27/17



UWAC J2 07/18/18



UWAC J2 06/21/19



UWAC J2 07/24/20



UWAC J2 07/13/21

UWAC T7 was restored in September 2014. GPS: 33.7526, -118.414563



UWAC T7 PRE-RESTORATION 08/14/14



UWAC T7 07/07/16



UWAC T7 07/27/17



UWAC T7 07/18/18



UWAC T7 06/21/19




UWAC T7 07/24/20



UWAC T7 07/13/21

Hawthorne



Hawthorne 2 this large pinnacle within Block 2 is easily found by divers and will serve as the starting point for video transects and photos of the site. The photos below show the pinnacle at heading 180 degrees and 90 degrees.

GPS: 33.75064, -118.416097



HAW 2 Heading 180 08/10/16






HAW 2 Heading 180 07/20/18



HAW 2 Heading 180 07/18/19



HAW 2 Heading 180 11/11/20



HAW 2 Heading 180 07/09/21



HAW 2 Heading 90 08/10/16



HAW 2 Heading 90 08/25/17



HAW 2 Heading 90 07/20/18



HAW 2 Heading 90 07/18/19




HAW 2 Heading 90 11/11/20



HAW 2 Heading 90 07/09/21

Point Fermin



Point Fermin, Block J7 (FERM J7) north-south running ridge has been well documented with video footage pre and post restoration. GPS: 33.703028, -118.290167



FERM J7 9/25/15



FERM J7 8/10/16



FERM J7 7/7/17



FERM J7 7/17/18



FERM J7 08/07/19




FERM J7 07/29/20



FERM J7 06/11/2021

White Point



White Point, Block 12. GPS: 33.71297, -118.3165. Video transect starts from the center of block 12 and goes 10-meters with a 0-degree heading. Then turns to a 90-degree heading and proceeds 30-meters.



White Point. Shallow depth urchin density conditions. 01/17/2019

White Point, Block 12 (WPT 12) east-west running ridge with large boulder directly 7-meters from the center of block 12 with a 0-degree heading. Established permanent photo plot. GPS: 33.71297, -118.3165



WPT 12 02/07/2020



WPT 12 07/24/2020



WPT 12 06/11/2021

2021 Video Transects (video files available by request)

Video transects are recorded annually at specific GPS points per site. Transect lines are drawn on the maps above for each site. Marguerite T10 video transect was discontinued in 2020 due to budgetary restraints, as well as proximity to T16 video transect which displays similar conditions.

Files

Honeymoon Cove:

1.0_HoneymoonCove_Videotransect_2021

Underwater Arch Cove:

2.0_UnderwaterArch_006_Videotransect_2021

2.1_UnderwaterArch_T7_Videotransect_2021

Marguerite:

3.0_Marguerite_T16_Videotransect_2021

Hawthorne:

4.0_Hawthorne_Videotransect_2021

Point Fermin:

5.0_PointFermin_Videotransect_2021

White Point:

6.0_Whitepoint_Videotransect_2021

Timelapse Videos of Sites (video files available by request)

Videos were taken at set blocks per site pre and post restoration. Each video consists of the same transect defined by GPS coordinates during summer months in different years.

Files

Honeymoon Cove:

1.1_Timelapse_HoneymoonCove_Videotransect_2021

Underwater Arch:

2.2_Timelapse_UnderwaterArch_Videotransect_2021

Marguerite:

3.1_Timelapse_Marguerite_Videotransect_2021

Hawthorne:

4.1_Timelapse_Hawthorne_Videotransect_2021

Point Fermin:

5.1_Timelapse_PointFermin_Videotransect_2021

White Point

6.1_Timelapse_Whitepoint_Videotransect_2021